

A QUANTITATIVE CHARACTERISATION OF FUNCTIONS WITH LOW AVILES GIGA ENERGY ON CONVEX DOMAINS

ANDREW LORENT

ABSTRACT. Given a connected Lipschitz domain Ω we let $\Lambda(\Omega)$ be the subset of functions in $W^{2,2}(\Omega)$ with $u = 0$ on $\partial\Omega$ and whose gradient (in the sense of trace) satisfies $\nabla u(x) \cdot \eta_x = 1$ where η_x is the inward pointing unit normal to $\partial\Omega$ at x . The functional $I_\epsilon(u) = \frac{1}{2} \int_\Omega \epsilon^{-1} |1 - |\nabla u|^2|^2 + \epsilon |\nabla^2 u|^2$ minimised over $\Lambda(\Omega)$ serves as a model in connection with problems in liquid crystals and thin film blisters, it is also the most natural higher order generalisation of the Modica Mortola functional. In [Ja-Ot-Pe 02] Jabin, Otto, Perthame characterised a class of functions which includes all limits of sequences $u_n \in \Lambda(\Omega)$ with $I_{\epsilon_n}(u_n) \rightarrow 0$ as $\epsilon_n \rightarrow 0$. A corollary to their work is that if there exists such a sequence (u_n) for a bounded domain Ω , then Ω must be a ball and (up to change of sign) $u := \lim_{n \rightarrow \infty} u_n = \text{dist}(\cdot, \partial\Omega)$. We prove a quantitative generalisation of this corollary for the class of bounded convex sets.

There exists positive constant γ_1 such that if Ω is a convex set of diameter 2 and $u \in \Lambda(\Omega)$ with $I_\epsilon(u) = \beta$ then $|B_1(x) \Delta \Omega| \leq c\beta^{\gamma_1}$ for some x and

$$\int_\Omega \left| \nabla u(z) + \frac{z-x}{|z-x|} \right|^2 dz \leq c\beta^{\gamma_1}.$$

A corollary of this result is that there exists positive constant $\gamma_2 < \gamma_1$ such that if Ω is convex with diameter 2 and C^2 boundary with curvature bounded by $\epsilon^{-\frac{1}{2}}$, then for any minimiser v of I_ϵ over $\Lambda(\Omega)$,

$$\|v - \zeta\|_{W^{1,2}(\Omega)} \leq c(\epsilon + \inf_y |\Omega \Delta B_1(y)|)^{\gamma_2}$$

where $\zeta(z) = \text{dist}(z, \partial\Omega)$. Neither of the constants γ_1 or γ_2 are optimal.

1. INTRODUCTION

We consider the following functional

$$I_\epsilon(u) = \frac{1}{2} \int_\Omega \epsilon^{-1} |1 - |\nabla u|^2|^2 + \epsilon |\nabla^2 u|^2 \quad (1)$$

the study of which arises from a number of sources, one of the earliest and most important is the article by Aviles, Giga [Av-Gi 87]. We will refer to the quantity $I_\epsilon(u)$ as the Aviles-Giga energy of function u . Functional I_ϵ is usually minimised over the space of functions $u \in W^{2,2}(\Omega)$ where $u(x) = 0$ and $\nabla u(x) \cdot \eta_x = 1$ on $\partial\Omega$ (in the sense of trace) where η_x is the inward pointing unit normal, we will denote this space of functions by $\Lambda(\Omega)$.

Aviles, Giga raised the problem of the study of the limiting behavior of I_ϵ as $\epsilon \rightarrow 0$ in connection with the theory of smectic liquid crystals [Av-Gi 87]. In [Gi-Or 97] Gioia, Ortiz studied I_ϵ as a model for thin film blisters. Jin, Kohn [Ji-Ko 00] introduced the by now classic method of estimating the energy by ‘divergence of vectorfields’. A related functional arising from micromagnetics was studied by Riviere, Serfaty [Ri-Se 01], in this case the functional acts on vector fields m (in two dimensions) satisfying $|m| = 1$ in Ω and the functional is given by

2000 *Mathematics Subject Classification.* 49N99.

Key words and phrases. Aviles Giga functional.

$M_\epsilon(m) = \epsilon \int_\Omega |\nabla m|^2 + \epsilon^{-1} \int_{\mathbb{R}^2} |\nabla^{-1} \operatorname{div} \tilde{m}|^2$ where $\tilde{m}$ is vectorfield m extended trivially by 0 outside Ω . For the Aviles Giga functional we minimise over curl free vector fields and the functional forces the norm of the vector field to be 1 with weighting ϵ^{-1} while constraining an ϵ multiple of the L^2 norm (squared) of the gradient, on the other hand the micromagnetics functional is minimised over vectorfields whose norm is taken to be 1 from the outset and the functional forces the vector field to be divergence free with weighting ϵ^{-1} while again constraining an ϵ multiple of the L^2 norm (squared) of the gradient. Functional M_ϵ is much more rigid and very much stronger results are known for it than for I_ϵ , see [Al-Ri-Se 02],[Ri-Se 01],[Am-Ki-Ri 02],[Am-Le-Ri 03].

Roughly speaking, the conjecture is that as $\epsilon \rightarrow 0$ the energy of minimisers of I_ϵ will converge to a collection of curves on which the gradient of the minimisers make a jump of order $o(1)$ perpendicularly across the curve. This has already been proved for functional M_ϵ [Ri-Se 01]. A way to think about this is the following, given a connected Lipschitz domain Ω let w be the distance from $\partial\Omega$ and let v_ϵ be w convolved by a convolution kernel of diameter ϵ , the regions where $|\nabla v_\epsilon| \not\sim 1$ will be exactly the ϵ neighborhoods of the curves on which ∇w has a jump discontinuity. If Ω is a ball ∇w will have a discontinuity only at one point, in all other cases there will be non trivial curves of singularities and for the specific function v_ϵ , it is exactly in an ϵ neighborhood of these curves that the energy will concentrate. The conjecture is that what we can observe directly for v_ϵ will hold true for the minimisers of I_ϵ .

The most natural way to study these questions is within the frame work of Γ convergence. One of the earliest successes of Γ convergence was the characterisation of the Γ limit of the so called Modica Mortola functional $A_\epsilon(w) = \int_\Omega \epsilon |\nabla w|^2 + \epsilon^{-1} |1 - |w|^2|^2$ which is minimised over scalar functions w satisfying an integral condition of the form $\int_\Omega w = 0$. It was shown by Modica, Mortola [Mo-Mo 00] (confirming a conjecture of DeGiorgi) that the Γ limit of A_ϵ is a constant multiple of the H^{n-1} measure of the jump set J_w minimised over the space of functions $w \in \{v \in BV : v \in \{1, -1\} \text{ a.e. and } \int v = 0\}$. Given the elementary inequality

$$\epsilon |\nabla w|^2 + \epsilon^{-1} |1 - |w|^2|^2 \geq |\nabla w| |1 - |w|^2| \quad (2)$$

we have that for any sequence (w_n) of equibounded A_{ϵ_n} energy (for some subsequence $\epsilon_n \rightarrow 0$) has a uniform L^1 control of $\nabla \left(w_n - \frac{w_n^3}{3}\right)$ and the measure we obtain as the limit of this L^1 sequence of gradients will naturally be supported on the jump set of the limiting function. In some sense the nature of the Γ limit of A_ϵ could be anticipated from (2).

Functional I_ϵ is the most natural higher order generalisation of A_ϵ , in the case of I_ϵ the conjectured Γ limit is surprising, this is part of the reason that functional I_ϵ has received so much attention. The first works on identifying the Γ limit are by Aviles, Giga [Av-Gi 87] and Jin, Kohn [Ji-Ko 00], later these ideas were developed by Ambrosio, DeLellis, Mantegazza [Am-De-Ma 99], roughly speaking the limiting function space is conjectured to have a structure similar to the space of functions whose gradient is BV and the limiting energy is conjectured to have the form $\int_{J_{\nabla u}} |\nabla u^+ - \nabla u^-|^3 dH^1$. Much progress has been made on this conjecture, particularly equi-coercivity of I_ϵ has been shown independently in [Am-De-Ma 99] and in the work of Desimone, Kohn, Muller, Otto [De-Ko-Mu-Ot 00]. A proposed limiting function space $AG(\Omega)$ and limiting functional I as been suggested in [Am-De-Ma 99] and it was shown that all limits of sequences of functions (u_n) with $\sup_n I_{\epsilon_n}(u_n) < \infty$ are such that $u_n \xrightarrow{W^{1,3}} u \in AG(\Omega)$ and $\liminf I_{\epsilon_n}(\nabla u_n) \geq I(u)$. The compactness proofs provided by [De-Ko-Mu-Ot 00] and [Am-De-Ma 99] are different but share some common ideas. The proof by [De-Ko-Mu-Ot 00]

¹the term $\int_{\mathbb{R}^2} |\nabla^{-1} \operatorname{div} m|^2$ is the L^2 norm of the Hodge projection onto curl free vector fields

identifies the set of functions Φ

$$\int |\nabla \Phi(\nabla u)| \leq c \int |\nabla^2 u| |1 - |\nabla u|^2| \text{ for any } C^2 \text{ function } u, \quad (3)$$

influenced by ideas of Tartar and Murat on compensated compactness [Ta 79] [Mu 78] the authors are able to prove that this set of Φ is sufficiently rich so as to force ∇u_n to converge strongly. In [Av-Gi 87] the authors (building on work of Jin Kohn [Ji-Ko 00]) found two third order polynomials Σ_1 and Σ_2 such that

$$\int |\operatorname{div} \Sigma_i(\nabla u)| \leq c \int |\nabla^2 u| |1 - |\nabla u|^2| \text{ for any } C^2 \text{ function } u, \text{ for } i = 1, 2. \quad (4)$$

Using some elementary and surprising identities satisfied by $\Sigma_1(\nabla u), \Sigma_2(\nabla u)$ a different approach to compactness was found. Rather naturally considering (4), the function space $AG(\Omega)$ proposed by [Am-De-Ma 99] is given by the set of functions v for which $\operatorname{div}(\Sigma_i(\nabla v))$ forms a Radon measure for $i = 1, 2$ and the limiting energy functional $I(v)$ is given by the total absolute value of this measure on Ω .

Given vector field w let $\chi(\xi, w) := \mathbb{1}_{\{\xi \cdot w > 0\}}$, Jabin, Perthame [Ja-Pe 97] showed that gradients of sequences of bounded Aviles-Giga energy (in fact their method extends to more general functionals) are compact and the limit ∇u satisfies a kinetic equation of the form $\xi \cdot \nabla_x \chi(\xi, R(\nabla u)) = q$ where q is the distribution derivative with respect to ξ of some measure on $\mathbb{R}_\xi^2 \times \mathbb{R}_x^2$ and R is the rotation given by $R(x, y) = (-y, x)$. By application of kinetic averaging lemmas [Di-Li-Me 91] this leads to some regularity; $\nabla u \in W^{s, q}$ for all $0 \leq s < \frac{1}{5}$, $q < \frac{5}{3}$ and using the kinetic equation a different proof of compactness was found. The kinetic equation deduced by [Ja-Pe 97] was motivated by the characterisation of the set of Φ satisfying (3) given in [De-Ko-Mu-Ot 00], indeed defining $\tilde{\Phi}(z) = |z|^2 e$ for $z \cdot e > 0$ and 0 otherwise, in [De-Ko-Mu-Ot 00] it was shown that a sequence Φ_n satisfying (2) could be found that approximates $\tilde{\Phi}$ pointwise. Using the kinetic equation deduced in [Ja-Pe 97], Jabin, Otto, Perthame [Ja-Ot-Pe 02] were able to characterise zero energy limits (and the domains that allow them) for I_ϵ , in fact their result is stronger, they showed that if a divergence free vector field m satisfies the kinetic equation $\xi \cdot \nabla \chi(m, \xi) = 0$, $|m(x)| = 1$ a.e. in Ω and $m(x) \cdot \eta_x = 0$ on $\partial\Omega$ then either Ω is a strip and m is a constant or $\Omega = B_r(x)$ for some $r > 0$, $x \in \mathbb{R}^2$ and $m(z) = \left(\frac{z-x}{|z-x|}\right)^\perp$ or $m(z) = -\left(\frac{z-x}{|z-x|}\right)^\perp$. An analogous result for zero energy limits of M_ϵ is stated in [Le-Ri 02] and is a consequence of the main theorem of [Am-Le-Ri 03].

As a corollary, given a sequence $u_n \in \Lambda(\Omega)$ and $\epsilon_n \rightarrow 0$ such that $I_{\epsilon_n}(u_n) \rightarrow 0$ as $n \rightarrow \infty$, letting u be the limit of this sequence, the vector field $R(\nabla u)$ satisfies the hypothesis stated and hence we have (up to a sign) a complete description of ∇u .

The main theorem of this paper is a quantitative generalisation of the corollary to Jabin, Otto, Perthame theorem over the class of bounded convex sets.

Theorem 1. *Let $\epsilon > 0$ and Ω be a convex domain with diameter 2. Let $u \in W^{2,2}(\Omega)$ with $u = 0$ on $\partial\Omega$ and $\nabla u(x) \cdot \eta_x = 1$ of $\partial\Omega$ (in the sense of trace) where η_x is the inward pointing unit normal. Then there exists positive constants $\mathcal{C} > 1$ and $\gamma < 1$ such that for some $x \in \Omega$,*

$$|\Omega \Delta B_1(x)| \leq \mathcal{C} (I_\epsilon(u))^\gamma$$

and

$$\int_\Omega \left| \nabla u(z) + \frac{z-x}{|z-x|} \right|^2 dz \leq \mathcal{C} (I_\epsilon(u))^\gamma.$$

Corollary 1. *Let Ω be a convex set with diameter 2, C^2 boundary and curvature bounded above by $\epsilon^{-\frac{1}{2}}$. Let $\Lambda(\Omega) := \{u \in W^{2,2}(\Omega) : u = 0 \text{ on } \partial\Omega \text{ and } \nabla u(z) \cdot \eta_z = 1 \text{ for } z \in \partial\Omega\}$. There exists positive constants $\mathcal{C} > 1$ and $\lambda < 1$ such that if u is a minimiser of I_ϵ over $\Lambda(\Omega)$, then*

$$\|u - \zeta\|_{W^{1,2}(\Omega)} \leq \mathcal{C} \left(\epsilon + \inf_y |\Omega \Delta B_1(y)| \right)^\lambda \quad (5)$$

where $\zeta(z) = \text{dist}(z, \partial\Omega)$.

In Theorem 1 we take $\gamma = 512^{-1}$ and in Corollary 1, $\lambda = 3000^{-1}$. Neither constant is optimal.

1.1. Background. Given a sequence $\epsilon_n \rightarrow 0$ and $u_n \in \Lambda(\Omega)$ with $\limsup I_{\epsilon_n}(u_n) < \infty$, let u be the limit of u_n , the vector valued measure given by $\nu_u := (\text{div}\Sigma_1(\nabla u), \text{div}\Sigma_2(\nabla u))$ (where Σ_1, Σ_2 are the third order polynomials that satisfy (4)) gives us the expression of the limiting energy, i.e. $I(u) = \|\nu_u\|(\Omega)$. If we consider the 1-dimensional part of the measure

$$\Gamma := \left\{ x : \limsup_{r \rightarrow 0} \frac{\|\nu_u(B_r(x))\|}{r} > 0 \right\}$$

it has been shown that Γ is 1-rectifiable [De-Ot 03] (see also [De-Ot-We 03]) and an analogous result has been shown for M_ϵ [Am-Ki-Ri 02]. It was also shown ∇u has jump discontinuities across the rectifiable set Γ exactly as would be the case if ∇u was BV and its jump set was given by Γ . However it is not known (even if u_n are the minimisers of I_{ϵ_n}) if measure $\|\nu_u\|$ is even singular with respect to Lebesgue measure. Note that for the function M_ϵ the minimiser of the limiting energy is known to be rectifiable [Am-Le-Ri 03], for a sequence with only equibounded energy the measure is not known to be singular.

The original motivation for Theorem 1 was to prove a version of it for $\Omega = B_1(0)$ without boundary conditions, under the hypotheses $\int_{B_1} |1 - |\nabla u|^2| |\nabla^2 u| = \beta$, $\int_{B_1} |1 - |\nabla u|^2| \leq \epsilon$ and $\sup \{\|u - A\|_{L^\infty(B_1(0))} : A \text{ is affine with } |\nabla A| = 1\} \leq 1000^{-1}$, the conclusion in this case would be that there exists a smooth function ψ with $|\nabla \psi| = 1$ everywhere such that $\|\nabla u - \nabla \psi\|_{L^2(B_{2^{-1}}(0))} \leq c\beta^\gamma$ for some $\gamma > 0$. This is a kind of quantitative version of the main proposition required to prove compactness in [Am-De-Ma 99], (see Proposition 4.6). The hope is to use such a quantitative result to show $\|\nu_u\|$ is singular, or at least that ∇u is continuous at H^1 a.e. point outside Γ , we will address these issues in a forthcoming paper [Lo pr].

The many strong results about measure $\|\nu_u\|$ (and the measure that gives the limiting functional for the micromagnetics function) have been achieved by characterising various kinds of *blow up* of the measure and understanding well the absolute (i.e. non quantitative) situation in the limit [Am-Ki-Ri 02], [De-Ot 03], [De-Ot-We 03], [Ja-Ot-Pe 02], [Am-Le-Ri 03]. In some sense there are only two possibilities, to take a limit and have an absolute situation and to understand the measure from this, or to stop before the limit and have a non-absolute situation and try and understand something about it with a quantitative theorem. Our primary motivation in proving a quantitative version of Jabin-Otto-Perthame Theorem was so as to obtain a result that could be used for the latter approach.

By Poincaré's inequality it is easy to see $\inf_{\Lambda(\Omega)} I_\epsilon \geq c\epsilon$ and so Theorem 1 follows from the following slightly more general result.

Theorem 2. *Let Ω be a convex body centered on 0 with $\text{diam}(\Omega) = 2$. Let $\beta > 0$, suppose $u : W^{2,2}(\Omega) \rightarrow \mathbb{R}$ is a function satisfying*

$$\int_{\Omega} |1 - |\nabla u|^2| |\nabla^2 u| dz \leq \beta \quad (6)$$

and

$$\int_{\Omega} \left| 1 - |\nabla u|^2 \right|^2 dz \leq \beta^2 \quad (7)$$

and in addition u satisfies $u = 0$ on $\partial\Omega$ and $\nabla u(z) \cdot \eta_z = 1$ on $\partial\Omega$ in the sense of trace where η_z is the inward pointing unit normal to $\partial\Omega$ at z .

Then there exists positive constant $C_1 > 0$ such that $|B_1(0) \triangle \Omega| < C_1 \beta^{\frac{1}{256}}$ and

$$\int_{\Omega} \left| \nabla u(z) + \frac{z}{|z|} \right|^2 dz \leq C_1 \beta^{\frac{1}{256}}. \quad (8)$$

Acknowledgments. The larger part of this paper was written while the author was the Emma e Giovanni Sansone Junior Visitor at Centro di Ricerca Matematica Ennio De Giorgi, Pisa. The hospitality and support this institute is gratefully acknowledged.

2. SKETCH OF THE PROOF

2.1. Sketch of the proof of Theorem 2. While the proof for convex domains is slightly involved, there are only a couple of ideas that are really central. We will sketch the proof for the case $\Omega = B_1(0)$, ignoring (without comment) many technicalities in order to give an impression of the basic skeleton.

The real engine of the proof is the characterisation in [De-Ko-Mu-Ot 00] of the set of Φ such that (3) is satisfied. As mentioned in the introduction, as consequence of the characterisation it was shown there exists a sequence of Φ_n satisfying (3) that converge pointwise to the function $\tilde{\Phi}(z) = |z|^2 e$ for $z \cdot e > 0$ and 0 otherwise. Following closely the proof of this it is possible to extract the existence of functions Φ_θ and Ψ_θ with $\|\nabla \Phi_\theta\| \leq c\beta^{-\frac{1}{4}}$, $\|\Psi_\theta\| \leq c\beta^{-\frac{1}{4}}$, $\|\nabla \Psi_\theta\| \leq c\beta^{-\frac{1}{2}}$ such that the following two inequalities hold.

Let $\Lambda_\theta(z) := \theta$ for $z \cdot \theta > 0$ and 0 otherwise,

$$|\Phi_\theta(z) - \Lambda_\theta(z)| \leq c\beta^{\frac{1}{4}} \text{ for } z \in N_{\sqrt{\beta}}(S^1) \setminus B_{2\beta^{\frac{1}{4}}}(\theta) \quad (9)$$

and (letting $R(z_1, z_2) = (-z_2, z_1)$ be the anti-clockwise rotation)

$$\operatorname{div} \left[\Phi_\theta(R(\nabla w)) - \Psi_\theta(R(\nabla w)) \left(1 - |R(\nabla w)|^2 \right) \right] \leq c\beta^{-\frac{1}{2}} \left| 1 - |\nabla w|^2 \right| |\nabla^2 w| \text{ for any } w \in W^{2,1}. \quad (10)$$

Recall, for simplicity we have taken $\Omega = B_1(0)$, as $\nabla u(z) = -\frac{z}{|z|}$ on $\partial B_1(0)$ then we can extend u to a function $\tilde{u} : B_{11/10}(0) \rightarrow \mathbb{R}$ such that

$$\int_{B_{11/10}(0)} \left| 1 - |\nabla \tilde{u}|^2 \right| |\nabla^2 \tilde{u}| \leq c\beta, \quad \int_{B_{11/10}(0)} \left| 1 - |\nabla \tilde{u}|^2 \right|^2 \leq c\beta^2$$

and

$$\nabla \tilde{u}(z) = -\frac{z}{|z|} \text{ for any } z \in B_{11/10}(0). \quad (11)$$

It is more convenient to work with vectorfields that are *almost* curl free instead of *almost* divergence free. So notice that (9) can be rewritten as

$$|R(\Phi_\theta(z)) - R(\Lambda_\theta(z))| \leq c\beta^{\frac{1}{4}} \text{ for } z \in N_{\sqrt{\beta}}(S^1) \setminus B_{2\beta^{\frac{1}{4}}}(\theta) \quad (12)$$

and we have $\int_{B_{11/10}(0)} \left| \operatorname{curl} \left[R(\Phi_\theta(R(\nabla \tilde{u}))) - R(\Psi_\theta(R(\nabla \tilde{u}))) \left(1 - |\nabla \tilde{u}|^2 \right) \right] \right| \leq c\sqrt{\beta}$. By the quantitative Hodge decomposition type theorem from [Am-De-Ma 99] (Theorem 4.3) we can find a scalar valued function w_θ such that

$$\int_{B_{11/10}(0)} \left| \nabla w_\theta - \left(R(\Phi_\theta(R(\nabla \tilde{u}))) - R(\Psi_\theta(R(\nabla \tilde{u}))) \left(1 - |\nabla \tilde{u}|^2 \right) \right) \right| \leq c\sqrt{\beta}. \quad (13)$$

The real power of (13) is that on the annulus $\mathcal{A} := B_{11/10}(0) \setminus B_1(0)$ we know that $\nabla \tilde{u}(z) = -\frac{\bar{z}}{|z|}$ and hence given inequality (12) (and the fact that $|\nabla \tilde{u}| = 1$ on $\mathcal{A}$) we have a that $\Phi_\theta(R(\nabla \tilde{u}(z))) \in N_{\beta^{\frac{1}{4}}}(\theta)$ for any $z \in \mathcal{A} \cap H(R\theta, 0)$, see figure 1.




FIGURE 1.

In much the same way in the ball $B_1(0)$, by inequalities (12), (13) and $\int_{B_1(0)} |1 - |\nabla \tilde{u}|^2|^2 \leq \beta^2$ we have that there exists a large set $\mathcal{G} \subset B_1(0) \cap H(0, R\theta)$, with $|B_1(0) \setminus \mathcal{G}| \leq \sqrt{\beta}$ such that if $z \in \mathcal{G}$ then $\nabla w_\theta(z) \in B_{\beta^{\frac{1}{4}}}(R\theta)$ or $\nabla w_\theta(z) \in B_{\beta^{\frac{1}{4}}}(0)$ depending on whether $R(\nabla u(z)) \cdot \theta > 0$ or $R(\nabla u(z)) \cdot \theta \leq 0$.

It is not hard to see we can find points $a, b \in N_{\beta^{\frac{1}{8}}}(\langle \theta \rangle \cap \partial B_1(0))$ with $|a - b| \sim 2 \cdot \theta \cdot \frac{b-a}{|b-a|} > 0$, the angle between $\frac{b-a}{|b-a|}$ and θ is at least $\beta^{\frac{1}{8}}$ and $H^1([a, b] \setminus \mathcal{G}) \leq \beta^{\frac{1}{4}}$. Let $\mathcal{G}_1 = \{x \in \mathcal{G} : \nabla u(z) \cdot R^{-1}(\theta) > 0\}$ and $\mathcal{G}_2 = \mathcal{G} \setminus \mathcal{G}_1$. As can be seen from figure 1 we can connect a to b with a path $\Gamma \subset \mathcal{A}$ so

$$\begin{aligned} |w_\theta(b) - w_\theta(a)| &= \left| \int_\Gamma \nabla w_\theta(z) t_z dH^1 z \right| \geq \left| R\theta \cdot \left(\int_\Gamma t_z dH^1 z \right) \right| - c\beta^{\frac{1}{4}} \\ &= \left| R\theta \cdot \frac{b-a}{|b-a|} \right| |b-a| - c\beta^{\frac{1}{4}}. \end{aligned} \tag{14}$$

On the other hand

$$\begin{aligned}
|w_\theta(b) - w_\theta(a)| &= \left| \int_{[a,b]} \nabla w_\theta(z) \frac{b-a}{|b-a|} dH^1 z \right| \leq \left| \int_{[a,b] \cap \mathcal{G}_1} \nabla w_\theta(z) \frac{b-a}{|b-a|} dH^1 z \right| + c\beta^{\frac{1}{4}} \\
&\leq \left| \int_{[a,b] \cap \mathcal{G}_1} R\theta \cdot \frac{b-a}{|b-a|} dH^1 z \right| + c\beta^{\frac{1}{4}} \\
&= \left| R\theta \cdot \frac{b-a}{|b-a|} \right| H^1([a,b] \cap \mathcal{G}_1) + c\beta^{\frac{1}{4}}
\end{aligned} \tag{15}$$

and since $\left| R\theta \cdot \frac{b-a}{|b-a|} \right| \geq \beta^{\frac{1}{8}}$ so putting (14) and (15) together

$$|a-b| \leq H^1([a,b] \cap \mathcal{G}_1) + \frac{c\beta^{\frac{1}{4}}}{\left| R\theta \cdot \frac{b-a}{|b-a|} \right|} \leq H^1([a,b] \cap \mathcal{G}_1) + c\beta^{\frac{1}{8}}.$$

So by arguing in the same way for lines parallel to $[a,b]$ by Fubini's theorem we can show $\left| H\left(\frac{a+b}{2}, R\left(\frac{b-a}{|b-a|}\right)\right) \setminus \mathcal{G}_1 \right| \leq c\beta^{\frac{1}{8}}$. Thus all but $\beta^{\frac{1}{8}}$ points $z \in B_1(0) \cap H(0, R(\theta))$ are such that $\nabla u(z) \cdot R^{-1}(\theta) > 0$. As θ is arbitrary we can rephrase this the following way. Given $\phi \in S^1$ for all but $\beta^{\frac{1}{8}}$ points $z \in B_1(0) \cap H(0, \phi)$ are such that $\nabla u(z) \cdot (-\phi) > 0$.

Now take $\psi = \begin{pmatrix} \cos \beta^{\frac{1}{16}} \\ \sin \beta^{\frac{1}{16}} \end{pmatrix}$. For all but $\beta^{\frac{1}{8}}$ points in $H(0, e_1) \cap H(0, -\psi) \cap H(0, -e_2)$ we have that $\nabla u(z) \cdot (-e_1) > 0$ and $\nabla u(z) \cdot \psi > 0$, it is not hard to show this implies $|\nabla u(z) \cdot e_1| \leq c\beta^{\frac{1}{16}}$ and since $\nabla u(z) \cdot e_2 > 0$ and $|\nabla u(z)| \sim 1$ we have $\nabla u(z) \in B_{c\beta^{\frac{1}{16}}}(e_2)$ with an exceptional set of measure less than $c\beta^{\frac{1}{8}}$. So integrating a carefully chosen line inside $H(0, e_1) \cap H(0, -\psi) \cap H(0, -e_2)$ and using the fact that $u = 0$ on $\partial B_1(0)$ we can show $|u(0) - 1| \leq c\beta^{\frac{1}{16}}$.

Now since $|\nabla u|$ is mostly very close to 1 and we have zero boundary condition, so avoiding technicalities assuming the coarea formula we have $\int_{\theta \in S^1} \int_{\mathbb{R}_+ \theta \cap B_1(0)} \left| |\nabla u(z)|^2 - 1 \right| dH^1 z dH^1 \theta \leq c\sqrt{\beta}$. Note also that for any $\theta \in S^1$, $u(\theta) = 0$ so by the fundamental theorem of Calculus

$$\begin{aligned}
\left| \int_{\mathbb{R}_+ \theta \cap B_1(0)} \nabla u(z) \cdot (-\theta) dH^1 z - 1 \right| &\leq |(u(0) - u(\theta)) - 1| \\
&\leq c\beta^{\frac{1}{16}}
\end{aligned}$$

so

$$\begin{aligned}
&\int_{\theta \in S^1} \int_{\mathbb{R}_+ \theta \cap B_1(0)} |\nabla u(z) + \theta|^2 dH^1 z dH^1 \theta \\
&= \int_{\theta \in S^1} \int_{\mathbb{R}_+ \theta \cap B_1(0)} |\nabla u(z)|^2 + 2\nabla u(z) \cdot \theta + |\theta|^2 dH^1 z dH^1 \theta \\
&\leq c\beta^{\frac{1}{16}}.
\end{aligned} \tag{16}$$

This concludes the sketch of the proof of Theorem 2.

2.2. Sketch of the proof of Corollary 1. In order to deduce Corollary 1 we need to apply Theorem 1 to the minimizer of I_ϵ over $\Lambda(\Omega)$. We can only do this if the minimiser has small energy (and from Theorem 1 we know it can only have small energy if Ω is close to a ball). For this reason it is necessary to construct a function in $\Lambda(\Omega)$ with this property. It turns out this is a surprisingly delicate task, it is achieved in Section 4 of the paper.

The obvious way to attempt the construction is to make some adaption of the function $\zeta(z) = \text{dist}(z, \partial\Omega)$, this function clearly satisfies the correct boundary condition. The first problem is

that $\nabla\zeta$ will have its gradient in BV and it is easy to construct examples of convex domains that are close to balls for which the singular part of $\nabla\zeta$ is widely spread over the domain. So it is necessary to convolve ζ , however convolution will destroy the boundary condition. To circumvent this obstacle, in a neighborhood of the boundary we convolve the ζ with a convolution kernel whose support decreases proportionally to the distance to the boundary. We make the assumption that $\partial\Omega$ is C^2 with curvature bounded above by $\epsilon^{-\frac{1}{2}}$ and this allows us to estimate the various error terms involved in differentiating a function convolved with a kernel of varying support.

3. PROOF OF THEOREM

It should be re-emphasized that the main calculations that makes this lemma work (specifically equation (23)) are very minor adoptions of the calculations in [De-Ko-Mu-Ot 00].

Lemma 1. *Let Ω be a convex body centered on 0 with $\text{diam}(\Omega) \leq 2$. Suppose $u : W^{2,1}(\Omega) \rightarrow \mathbb{R}$ satisfies (6) and (7). For each $\theta \in S^1$ define $\Lambda_\theta : \mathbb{R}^2 \rightarrow S^1$ be defined by*

$$\Lambda_\theta(z) = \begin{cases} \theta & \text{if } z \cdot \theta > 0, \\ 0 & \text{if } z \cdot \theta \leq 0. \end{cases} \quad (17)$$

Let $R \in SO(2)$ be the anti-clockwise rotation defined by $R(z_1, z_2) = (-z_2, z_1)$ and let $m = R(\nabla u)$, we will show there exists a set $\Gamma \subset S^1$ with $H^1(S^1 \setminus \Gamma) \leq c\beta^{\frac{1}{8}}$ and $-\Gamma = \Gamma$ such that for any $\theta \in \Gamma$ we can find function $w_\theta : \Omega \rightarrow \mathbb{R}$ with the property

$$\int_{\Omega} |\nabla w_\theta - R(\Lambda_\theta(m))| \leq c\beta^{\frac{1}{8}}. \quad (18)$$

Proof of Lemma 1. Let $M = 2 \left\lceil \frac{\beta^{-\frac{1}{4}}}{8} \right\rceil$, we divide S^1 into M disjoint connected subsets of length $\frac{2\pi}{M}$, denote them $A_1, A_2, \dots, A_M$. We assume they have been ordered sequentially, i.e. $\overline{A_i} \cap \overline{A_{i+1}} \neq \emptyset$ for $i = 1, 2, \dots, M-1$. Also assume they have been ordered so that $-A_i = A_{i+\frac{M}{2}}$ for $i = 1, 2, \dots, \frac{M}{2}$. Let

$$\mathcal{B} = \left\{ k \in \left\{ 1, 2, \dots, \frac{M}{2} \right\} : \left| \left\{ x \in \Omega : \frac{\nabla u(x)}{|\nabla u(x)|} \in \overline{A_k} \cup \overline{A_{k+\frac{M}{2}}} \right\} \right| \geq \beta^{\frac{1}{8}} \right\}.$$

Since $\text{Card}(\mathcal{B})\beta^{\frac{1}{8}} \leq |\Omega| \leq 4\pi$ we have that $\text{Card}(\mathcal{B}) \leq 4\pi\beta^{-\frac{1}{8}}$.

Let $\mathcal{D} := \{k \in \{2, 3, \dots, \frac{M}{2} - 1\} : \{k-1, k, k+1\} \cap \mathcal{B} \neq \emptyset\}$. A simple covering argument shows that $\text{Card}(\mathcal{D}) \leq c\beta^{-\frac{1}{8}}$. Let $\Gamma = \bigcup_{k \in \{2, 3, \dots, \frac{M}{2} - 1\} \setminus \mathcal{D}} \overline{A_k} \cup \overline{A_{k+\frac{M}{2}}}$. Note that for any $\theta \in \Gamma$ we have

$$\left| \left\{ x \in \Omega : \frac{\nabla u(x)}{|\nabla u(x)|} \in B_{2\beta^{\frac{1}{8}}}(\theta) \cup B_{2\beta^{\frac{1}{8}}}(-\theta) \right\} \right| \leq 3\beta^{\frac{1}{8}}. \quad (19)$$

So pick $\theta \in \Gamma$ without loss of generality we can assume $\theta = e_1$. Let $s : \mathbb{R} \rightarrow \mathbb{R}_+$ be a smooth monotone function where $s(x) = 0$ if $x \leq 0$ and $s(x) = x$ if $x > \beta^{\frac{1}{4}}$ and $\|\nabla^2 s\|_{L^\infty} \leq \beta^{-\frac{1}{4}}$ and $\|\nabla^3 s\|_{L^\infty} \leq \beta^{-\frac{1}{2}}$, it is clear such a function exists.

Let $\varphi(z) = s(z \cdot e_1) = s(z_1)$. Define $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$\begin{aligned} \Phi(z) &:= \varphi(z) \begin{pmatrix} z_1 \\ z_2 \end{pmatrix} + \left(\nabla \varphi(z) \cdot \begin{pmatrix} -z_2 \\ z_1 \end{pmatrix} \right) \begin{pmatrix} -z_2 \\ z_1 \end{pmatrix} \\ &= \begin{pmatrix} \varphi(z) z_1 + z_2^2 \varphi_{,1}(z) \\ \varphi(z) z_2 - z_2 z_1 \varphi_{,1}(z) \end{pmatrix}. \end{aligned} \quad (20)$$

Define

$$\Psi(z) = \begin{pmatrix} \Psi_1(z) \\ \Psi_2(z) \end{pmatrix} := \begin{pmatrix} -\varphi_{,1}(z) \\ \frac{z_2}{2} \varphi_{,11}(z) \end{pmatrix}. \quad (21)$$

Recall $m(z) := R(\nabla u(z))$ so m is divergence free. Note (using the fact $\varphi_{,2} \equiv 0$ and $\varphi_{,12} \equiv 0$ and $\operatorname{div} m \equiv 0$ for the third inequality, and using $\operatorname{div} m = 0$ for the last inequality)

$$\begin{aligned}
\operatorname{div} [\Phi(m)] &= \operatorname{div} \begin{pmatrix} \varphi(m)m_1 + m_2^2\varphi_{,1}(m) \\ \varphi(m)m_2 - m_2m_1\varphi_{,1}(m) \end{pmatrix} \\
&= (\varphi_{,1}(m)m_{1,1} + \varphi_{,2}(m)m_{2,1})m_1 + \varphi(m)m_{1,1} + 2m_2m_{2,1}\varphi_{,1}(m) \\
&\quad + m_2^2(\varphi_{,11}(m)m_{1,1} + \varphi_{,12}(m)m_{2,1}) + (\varphi_{,1}(m)m_{1,2} + \varphi_{,2}(m)m_{2,2})m_2 \\
&\quad + \varphi(m)m_{2,2} - ((m_{1,2}m_2 + m_1m_{2,2})\varphi_{,1}(m) \\
&\quad + m_1m_2(\varphi_{,11}(m)m_{1,2} + \varphi_{,12}(m)m_{2,2})) \\
&= m_1\varphi_{,1}(m)m_{1,1} + 2m_2m_{2,1}\varphi_{,1}(m) + m_2^2m_{1,1}\varphi_{,11}(m) + m_2m_{1,2}\varphi_{,1}(m) \\
&\quad - ((m_{1,2}m_2 + m_1m_{2,2})\varphi_{,1}(m) + m_1m_2m_{1,2}\varphi_{,11}(m)) \\
&= 2\varphi_{,1}(m)(m_1m_{1,1} + m_2m_{2,1}) - \varphi_{,11}(m)m_2(m_1m_{1,2} + m_2m_{2,2}). \tag{22}
\end{aligned}$$

Note also that

$$\begin{aligned}
\Psi(m) \cdot \nabla(1 - |m|^2) &= -\Psi(m) \cdot \begin{pmatrix} 2(m_1m_{1,1} + m_2m_{2,1}) \\ 2(m_1m_{1,2} + m_2m_{2,2}) \end{pmatrix} \\
&= 2\varphi_{,1}(m)(m_1m_{1,1} + m_2m_{2,1}) - m_2\varphi_{,11}(m)(m_1m_{1,2} + m_2m_{2,2})
\end{aligned}$$

so by (22) we have

$$\operatorname{div} [\Phi(m)] = \Psi(m) \cdot \nabla(1 - |m|^2). \tag{23}$$

Let $\tilde{\Phi} := R(\Phi)$ and $\tilde{\Psi} := R(\Psi)$ note $\operatorname{curl} [\tilde{\Phi}(m)] \stackrel{(23)}{=} \operatorname{div} [\Phi(m)] = \Psi(m) \cdot \nabla(1 - |m|^2)$. So

$$\begin{aligned}
\operatorname{curl} [\tilde{\Psi}(m)(1 - |m|^2)] &= \operatorname{div} [\Psi(m)](1 - |m|^2) + \Psi(m) \cdot \nabla(1 - |m|^2) \\
&= \operatorname{div} [\Psi(m)](1 - |m|^2) + \operatorname{curl} [\tilde{\Phi}(m)]. \tag{24}
\end{aligned}$$

Thus using the fact that $|\nabla \Psi(z)| \leq c|z| \|\nabla^3 \varphi\|_{L^\infty(\mathbb{R}^2)} \leq c\beta^{-\frac{1}{2}}|z|$ we have

$$\begin{aligned}
\operatorname{curl} [\tilde{\Phi}(m) - \tilde{\Psi}(m)(1 - |m|^2)] &\stackrel{(24)}{=} -\operatorname{div} [\Psi(m)](1 - |m|^2) \\
&= -(\Psi_{1,1}(m)m_{1,1} + \Psi_{1,2}(m)m_{2,1} + \Psi_{2,1}(m)m_{1,2} + \Psi_{2,2}(m)m_{2,2})(1 - |m|^2) \\
&\leq c\beta^{-\frac{1}{2}}|m| \left| 1 - |m|^2 \right| |\nabla m|. \tag{25}
\end{aligned}$$

Hence

$$\int_{\Omega} \left| \operatorname{curl} [\tilde{\Phi}(m) - \tilde{\Psi}(m)(1 - |m|^2)] \right| \leq c\beta^{-\frac{1}{2}} \int_{\Omega} |m| \left| 1 - |m|^2 \right| |\nabla m|$$

So using (25), note that if x is such that $|m(x)| \geq 2$ then for $J(x) := |m(x)|^3$ we have $|\nabla J(x)| \leq c \left| 1 - |m|^2 \right| |\nabla m|$ and so

$$\int_{\{x: 2 \leq |m(x)| \leq 4\}} |\nabla J(x)| dx \leq c \int_{\Omega} \left| 1 - |m|^2 \right| |\nabla m| \leq c\beta$$

so applying the Co-area formula we know $\int_8^{64} H^1(J^{-1}(s)) ds \leq c\beta$ thus we must be able to find $t \in [8, 64]$ such that $H^1(J^{-1}(t)) \leq c\beta$.

Let

$$\mathcal{G} := \{x \in \Omega : J(x) < t\}. \tag{26}$$

We claim we can construct a C^1 Lipschitz vector field $\Theta = (\Theta_1^1, \Theta_2^1) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ with the property $\Theta(x) = \tilde{\Phi}(x) - \tilde{\Psi}(x)(1 - |x|^2)$ for $x \in B_{64}(0)$. This can be seen by applying Whitney's

extension theorem (Theorem 1, Section 6.5 [Ev-Ga 92]) in the following way. Let $d_1(x) = \nabla \left[(\tilde{\Phi}(x) - \tilde{\Psi}(x)(1 - |x|^2)) \cdot e_1 \right]$ for $x \in \overline{B_{64}(0)}$ and let $d_2(x) = d_1(\frac{64x}{|x|})$, define $d : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ to be equal to d_1 in $B_{64}(0)$ and equal to d_2 outside it. Notice that d and $(\tilde{\Phi}(x) - \tilde{\Psi}(x)(1 - |x|^2)) \cdot e_1$ satisfy the hypotheses of Whitney's extension theorem so there exists a Lipschitz C^1 function Θ_1 which agrees with $(\tilde{\Phi}(x) - \tilde{\Psi}(x)(1 - |x|^2)) \cdot e_1$ on $B_{64}(0)$. There existence of Θ_2 follows in the same way.

Define $w : \Omega \rightarrow \mathbb{R}$ by

$$w(x) = \begin{cases} \tilde{\Phi}(m) - \tilde{\Psi}(m)(1 - |m|^2) & \text{for } x \in \mathcal{G} \\ 0 & \text{for } x \in \Omega \setminus \mathcal{G} \end{cases} \quad (27)$$

So if $x \in \mathcal{G}$,

$$\begin{aligned} \operatorname{curl}(w) &= \operatorname{curl} \left(\tilde{\Phi}(m) - \tilde{\Psi}(m)(1 - |m|^2) \right) \\ &\stackrel{(25),(26)}{\leq} c\beta^{-\frac{1}{2}} |1 - |m|| |\nabla m|. \end{aligned} \quad (28)$$

So if $x \in \operatorname{int}(\Omega \setminus \mathcal{G})$, $\operatorname{curl} \left(\tilde{\Phi}(m) - \tilde{\Psi}(m)(1 - |m|^2) \right) = 0$.

Since $m \in W^{1,1}(\Omega)$ and Θ is C^1 Lipschitz so the vector field $\Theta(m)$ is BV by Theorem 3.94 [Am-Fu-Pa 00]. So by Theorem 3.83 [Am-Fu-Pa 00] we have that w is also BV and the singular part of ∇w , which we denote by $[\nabla w]_s$, is supported on $J^{-1}(t) \cap \Omega$ and as $|\tilde{\Phi}(m(x))| \leq c|m(x)|^2$ and $|\tilde{\Psi}(m(x))| \leq c\beta^{-\frac{1}{4}}|m(x)|$ we have that

$$\operatorname{ess\,sup}_{J^{-1}(t) \cap \Omega} \left| \tilde{\Phi}(m(x)) - \tilde{\Psi}(m(x))(1 - |m(x)|^2) \right| \leq c\beta^{-\frac{1}{4}}$$

and thus $\|[\nabla w]_s\|(S) \leq c\beta^{-\frac{1}{4}}H^1(J^{-1}(t) \cap \Omega) \leq c\beta^{\frac{3}{4}}$. Now we know that for any set $S \subset \Omega$,

$$\|\operatorname{curl} w\|(S) \leq c\|\nabla w\|(S)$$

and so in particular

$$\|\operatorname{curl} w\|(J^{-1}(t)) \leq c\|\nabla w\|(J^{-1}(t)) \leq c\beta^{\frac{3}{4}}. \quad (29)$$

Thus

$$\begin{aligned} \|\operatorname{curl} w\|(\Omega) &\leq \|\operatorname{curl} w\|(J^{-1}(t)) + \|\operatorname{curl} w\|(\mathcal{G}) \\ &\quad + \|\operatorname{curl} w\|(\operatorname{int}(\Omega \setminus \mathcal{G})) \\ &\stackrel{(28),(29)}{\leq} c\beta^{\frac{3}{4}} + c\beta^{-\frac{1}{2}} \int_{\mathcal{G}} |1 - |m|| |\nabla m| \\ &\stackrel{(6)}{\leq} c\sqrt{\beta}. \end{aligned} \quad (30)$$

Now we try and understand the nature of vector field $\tilde{\Phi}(m(x)) - \tilde{\Psi}(m(x))(1 - |m(x)|^2)$. Note that if $z \in N_{\sqrt{\beta}}(S^1) \cap \{z_1 > 0\} \setminus \left(B_{2\beta^{\frac{1}{4}}}(e_2) \cup B_{2\beta^{\frac{1}{4}}}(-e_2) \right)$ then $\varphi(z) = z_1$, $\varphi_{,1}(z) = 1$ and so $\Phi(z) \stackrel{(20)}{=} \begin{pmatrix} z_1^2 + z_2^2 \\ 0 \end{pmatrix}$ on the other hand if $z \in N_{\sqrt{\beta}}(S^1) \cap \{z_1 \leq 0\} \setminus \left(B_{2\beta^{\frac{1}{4}}}(e_2) \cup B_{2\beta^{\frac{1}{4}}}(-e_2) \right)$ then $\varphi(z) = \varphi_{,1}(z) = 0$ and so $\Phi(z) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$.

Now if $z \in N_{\sqrt{\beta}}(S^1) \cap \{z_1 > 0\} \setminus (B_{2\beta^{\frac{1}{4}}}(e_2) \cup B_{2\beta^{\frac{1}{4}}}(-e_2))$ we have

$$\begin{aligned} \left| (\tilde{\Phi}(z) + \tilde{\Psi}(z)(1 - |z|^2)) - R(\Lambda_{e_1}(z)) \right| &\leq \left| \tilde{\Phi}(z) - R(\Lambda_{e_1}(z)) \right| + c\sqrt{\beta} \sup_{z \in N_{\sqrt{\beta}}(S^1)} \left| \tilde{\Psi}(z) \right| \\ &= \left| R \begin{pmatrix} z_1^2 + z_2^2 \\ 0 \end{pmatrix} - R \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right| + c\beta^{\frac{1}{4}} \\ &\leq c\beta^{\frac{1}{4}}. \end{aligned} \quad (31)$$

And if we have $z \in N_{\sqrt{\beta}}(S^1) \cap \{z_1 \leq 0\} \setminus (B_{2\beta^{\frac{1}{4}}}(e_2) \cup B_{2\beta^{\frac{1}{4}}}(-e_2))$ arguing in the same way we can conclude

$$\left| (\tilde{\Phi}(z) + \tilde{\Psi}(z)(1 - |z|^2)) - R(\Lambda_{e_1}(z)) \right| \leq c\beta^{\frac{1}{4}}. \quad (32)$$

Let $\Pi := \{z \in \Omega : |m(z)| \in (1 - \sqrt{\beta}, 1 + \sqrt{\beta})\}$ and let

$$\mathcal{E} := \left\{ x \in \Omega : \frac{\nabla u(x)}{|\nabla u(x)|} \in B_{2\beta^{\frac{1}{4}}}(e_2) \cup B_{2\beta^{\frac{1}{4}}}(-e_2) \right\}, \quad (33)$$

note from (19) we know $|\mathcal{E}| \leq 3\beta^{\frac{1}{8}}$. Note also $\sqrt{\beta}|\Omega \setminus \Pi| \leq c \int_{\Omega \setminus \Pi} |1 - |\nabla u|^2| \stackrel{(7)}{\leq} \beta$ thus

$$|\Omega \setminus \Pi| \leq c\sqrt{\beta}. \quad (34)$$

Now from (31) and (32)

$$\left| \int_{\Pi \setminus \mathcal{E}} (\tilde{\Phi}(m) - \tilde{\Psi}(m)(1 - |m|^2)) - R(\Lambda_{e_1}(m)) \right| \leq c\beta^{\frac{1}{4}} \quad (35)$$

on the other hand recalling the fact that $|\tilde{\Psi}(z)| \leq \beta^{-\frac{1}{4}}|z|$, $|\tilde{\Phi}(z)| \leq c|z|^2$ and using the definition of $\mathcal{G}$ (see (26)) we have

$$\begin{aligned} \left| \int_{\mathcal{G} \setminus \Pi} ((\tilde{\Phi}(m) - \tilde{\Psi}(m)(1 - |m|^2)) - R(\Lambda_{e_1}(m))) \right| \\ \leq c|\mathcal{G} \setminus \Pi| \\ \stackrel{(34)}{\leq} c\sqrt{\beta}. \end{aligned} \quad (36)$$

Thus applying (35) to (36) gives

$$\left| \int_{\mathcal{G} \setminus \mathcal{E}} ((\tilde{\Phi}(m) - \tilde{\Psi}(m)(1 - |m|^2)) - R(\Lambda_{e_1}(m))) \right| \leq c\beta^{\frac{1}{4}}. \quad (37)$$

Recall we have $|\mathcal{E}| \leq 3\beta^{\frac{1}{8}}$ so

$$\begin{aligned} \left| \int_{\mathcal{E} \cap \mathcal{G}} ((\tilde{\Phi}(m) - \tilde{\Psi}(m)(1 - |m|^2)) - R(\Lambda_{e_1}(m))) \right| &\leq c|\mathcal{E}| \\ &\leq c\beta^{\frac{1}{8}}. \end{aligned}$$

Putting this inequality together with (37) gives

$$\left| \int_{\mathcal{G}} ((\tilde{\Phi}(m) - \tilde{\Psi}(m)(1 - |m|^2)) - R(\Lambda_{e_1}(m))) \right| \leq c\beta^{\frac{1}{8}}. \quad (38)$$

So by definition of w (see (27)) we have that

$$\begin{aligned}
 \left| \int_{\Omega} w - R(\Lambda_{e_1}(m)) \right| &\stackrel{(38)}{\leq} c\beta^{\frac{1}{8}} + \left| \int_{\Omega \setminus \mathcal{G}} R(\Lambda_{e_1}(m)) \right| \\
 &\leq c\beta^{\frac{1}{8}} + |\Omega \setminus \mathcal{G}| \\
 &\stackrel{(34)}{\leq} c\beta^{\frac{1}{8}}.
 \end{aligned} \tag{39}$$

Now from (30) applying Theorem 4.3 from ([Am-De-Ma 99]) there exists $w_{e_1} \in W^{1,1}(\Omega)$ such that

$$\int_{\Omega} |\nabla w_{e_1} - w| \leq c\beta^{\frac{1}{8}} \tag{40}$$

thus putting this together with (39) and gives (18). $\square$

Lemma 2. *Let Ω be a convex body centered on 0 and let $u : W^{2,2}(\Omega) \rightarrow \mathbb{R}$ be a function satisfying (6) and (7) and $u = 0$ on $\partial\Omega$ and $\nabla u(z) \cdot \eta_z = 1$ on $\partial\Omega$ in the sense of trace, where η_z is the inward pointing unit normal to $\partial\Omega$ at z .*

For any $r > 0$ define $\Omega_r := N_r(\Omega)$, we will show we can construct a function $\tilde{u} : W^{2,1}(\Omega_r) \rightarrow \mathbb{R}$ satisfying

$$\int_{\Omega_r} |1 - |\nabla \tilde{u}|^2| |\nabla^2 \tilde{u}| \leq \beta, \quad \int_{\Omega_r} |1 - |\nabla \tilde{u}|^2| \leq \beta, \tag{41}$$

and

$$\tilde{u}(z) = \begin{cases} u(z) + r & \text{for } z \in \overline{\Omega} \\ r - d(z, \Omega) & \text{if } z \in \Omega_r \setminus \Omega \end{cases} \tag{42}$$

Proof of Lemma 2.

Step 1. We will show $\nabla u(x) = \eta_x$ for H^1 a.e. $x \in \partial\Omega$

Proof of Step 1. Recall $\nabla u \in W^{1,1}(\Omega)$ and ∇u is defined on $\partial\Omega$ in the sense of trace, as the trace operator is bounded we know $\int_{\partial\Omega} |\nabla u| dH^1 < \infty$.

We define

$$v(z) = \begin{cases} u(z) & \text{for } z \in \overline{\Omega} \\ 0 & \text{if } z \in \Omega_r \setminus \Omega \end{cases} \tag{43}$$

So note the vector field $\nabla v(z)$ is equal to $\nabla u(z)$ inside Ω and is zero outside, so by Theorem 3.8 [Am-Fu-Pa 00] $\nabla v \in BV(\Omega_r)$ and hence by Theorem 3.76 [Am-Fu-Pa 00] and Theorem 2, Section 5.3 [Ev-Ga 92] for H^1 a.e. $x \in \partial\Omega$ the following limits exist

$$\lim_{\rho \rightarrow 0} \int_{B^+(x, \eta_x)} |\nabla u(z) - \nabla u(x)| dz = 0 \tag{44}$$

and

$$\lim_{\rho \rightarrow 0} \int_{B^-(x, \eta_x)} |\nabla u(z)| dz = 0. \tag{45}$$

Let $w_x^\rho(z) = \frac{u(\rho(z-x))}{\rho}$, by (44) and (45) for any sequence $\rho_n \rightarrow 0$ we have $w_x^{\rho_n}(z) \xrightarrow{L^1} w_x$ as $n \rightarrow \infty$ where

$$w_x(z) = \begin{cases} \nabla u(x) \cdot z & \text{for } z \in H(0, \eta_x) \\ 0 & \text{for } z \in H(0, -\eta_x) \end{cases} \tag{46}$$

however w_x would not be curl free unless $\nabla u(x) = \lambda \eta_x$ for some $\lambda \in \mathbb{R}$. As we know $\nabla u(x) \cdot \eta_x = 1$ this implies $\nabla u(x) = \eta_x$ for H^1 a.e. $x \in \partial\Omega$. This completes the proof of Step 1.

Step 2. For any $z \in \Omega_r \setminus \Omega$, $\tilde{u}(z) = d(z, \partial\Omega_r)$.

Proof of Step 2. Note that $\|\nabla \tilde{u}\|_{L^\infty(\Omega_r \setminus \Omega)} \leq 1$. Let $x \in \partial\Omega_r$, let $q(x)$ be the metric projection onto a convex set Ω , i.e. the unique point for which $|x - q(x)| = d(x, \Omega)$. Since $x \in \partial\Omega_r = \partial(N_r(\Omega)) = \{x \in \Omega^c : d(x, \Omega) = r\}$ so $|x - q(x)| \geq r$, on the other hand we also know $d(x, \Omega) = r$ so there must exist $y \in \Omega$ such that $|x - y| < r + \delta$ for every $\delta > 0$, this implies $|x - q(x)| < r + \delta$ for every $\delta > 0$. Thus $|x - q(x)| = r$.

Since $\tilde{u}(x) = 0$ and $\tilde{u}(q(x)) = r$ and as $\tilde{u}$ is 1-Lipschitz on $\Omega_r \setminus \Omega$ this implies $\tilde{u}((1 - \alpha)x + \alpha q(x)) = \alpha r$ for any $\alpha \in [0, 1]$.

Now let $Q(z) := d(z, \partial\Omega_r)$. For every $x \in \partial\Omega_r$, $Q(q(x)) \leq |q(x) - x| = r$. As $\partial\Omega_r = \partial(N_r(\Omega))$ so we know $Q(q(x)) \geq r$ and thus have $Q(q(x)) = r$. We also know Q is 1-Lipschitz and $Q(x) = 0$, thus in the same way as before $Q((1 - \alpha)x + \alpha q(x)) = \alpha r$ for any $\alpha \in [0, 1]$. Therefor $Q(z) = \tilde{u}(z)$ for any $z \in [x, q(x)]$, $x \in \partial\Omega_r$ and this completes the proof of Step 1.

Step 3. We will show that $\tilde{u} \in W^{2,1}(\Omega_r)$ and that $\tilde{u}$ satisfies (41).

Proof of Step 3. First we claim that

$$\int_{\Omega_r \setminus \Omega} |\nabla^2 \tilde{u}| dz \leq c \text{ and hence } \nabla \tilde{u} \in W^{1,1}(\Omega_r \setminus \Omega). \quad (47)$$

Note $\tilde{u}|_{\Omega_r \setminus \Omega}$ is a 1-Lipschitz function with $\tilde{u} = 0$ on $\partial\Omega_r$ and $\tilde{u} = r$ on $\partial\Omega$. So by the Co-area formula we have $\int_{\Omega_r \setminus \Omega} |\nabla^2 \tilde{u}| dx = \int_0^r \int_{\tilde{u}^{-1}(h)} |\nabla^2 \tilde{u}| dH^1 dh$. Recall $\tilde{u}(x) = r - d(z, \Omega)$ for $z \in \Omega_r \setminus \Omega$, so as Ω is convex for every $z \in \Omega_r \setminus \Omega$ there is a unique point $b(z) \in \partial\Omega$ such that $d(z, \Omega) = |b(z) - z|$ and $\nabla \tilde{u}(z) = \frac{b(z) - z}{|b(z) - z|}$, since b is a continous function this shows that $\tilde{u}$ is C^1 in $\Omega_r \setminus \overline{\Omega}$ and $|\nabla \tilde{u}| = 1$ on $\Omega_r \setminus \overline{\Omega}$. Note this also implies $\tilde{u}^{-1}(h)$ is the boundary of a smooth convex set and for each $z \in \tilde{u}^{-1}(h)$, $\nabla \tilde{u}(z)$ is normal to the tangent of $\tilde{u}^{-1}(h)$ at z , so $\int_{\tilde{u}^{-1}(h)} |\nabla^2 \tilde{u}| dH^1 \leq c$ for all $h \in [0, r]$ and hence (47) is shown.

Since Ω is an extension domain by Theorem 1, Section 4.4 [Ev-Ga 92] there exists a function $p : W^{1,2}(\mathbb{R}^2) \rightarrow \mathbb{R}^2$ such that $p(z) = \nabla \tilde{u}(z)$ on Ω and $\text{Spt} p$ is compact. Similarly as $\Omega_r \setminus \Omega$ is an extension domain there exists a function $q : W^{1,1}(\mathbb{R}^2) \rightarrow \mathbb{R}^2$ such that $q(z) = \nabla \tilde{u}(z)$ on $\Omega_r \setminus \Omega$ and $\text{Spt} q$ is compact. We define $w : \Omega_r \rightarrow \mathbb{R}^2$ by $w := p\mathbb{1}_\Omega + q\mathbb{1}_{\Omega_r \setminus \Omega}$, by Theorem 3.83 [Am-Fu-Pa 00] $w \in BV(\Omega_r : \mathbb{R}^2)$ and since p and q agree on $\partial\Omega$ we have that ∇w as a measure is absolutely continuous with respect to Lebesgue measure (and hence $w \in W^{1,1}(\Omega_r : \mathbb{R}^2)$) and $\nabla w = \nabla p\mathbb{1}_\Omega + \nabla q\mathbb{1}_{\Omega_r \setminus \Omega}$. Now as $w = \nabla \tilde{u}$ a.e. in Ω_r we have that $\nabla \tilde{u} \in W^{1,1}(\Omega_r)$.

Since $\nabla^2 \tilde{u} \in L^1$ we know

$$\begin{aligned} \int_{\Omega_r} |1 - |\nabla \tilde{u}|^2| |\nabla^2 \tilde{u}| dz &= \int_{\Omega} |1 - |\nabla \tilde{u}|^2| |\nabla^2 \tilde{u}| dz + \int_{\Omega_r \setminus \Omega} |1 - |\nabla \tilde{u}|^2| |\nabla^2 \tilde{u}| dz \\ &= \int_{\Omega} |1 - |\nabla \tilde{u}|^2| |\nabla^2 \tilde{u}| dz \\ &\leq \beta. \end{aligned}$$

Similarly $\int_{\Omega_r} |1 - |\nabla \tilde{u}|^2| dz = \int_{\Omega} |1 - |\nabla \tilde{u}|^2| dz \leq \beta$. $\square$

Lemma 3. Let Ω be a convex body with $\text{diam}(\Omega) = 2$. Let $u : W^{2,2}(\Omega) \rightarrow \mathbb{R}$ be a function satisfying (6) and (7) and $u = 0$ on $\partial\Omega$ and $\nabla u(z) \cdot \eta_z = 1$ on $\partial\Omega$ in the sense of trace where η_z is the inward pointing unit normal to $\partial\Omega$ at z . For any $x, v \in \mathbb{R}^2$ let $H(x, v) := \{z \in \mathbb{R}^2 : (z - x) \cdot v > 0\}$.

Let $\Gamma \subset S^1$ be the set constructed in Lemma 1. Let $\mathcal{U} := \Omega_{1/10}$ be the convex body and $\tilde{u} : W^{2,1}(\mathcal{U}) \rightarrow \mathbb{R}$ be the function constructed in Lemma 2. Let R be the anti-clockwise rotation defined by $R(z_1, z_2) = (-z_2, z_1)$. Let $R_0 \in \{R^{-1}, R\}$.

For a.e. $\theta \in \Gamma$ there exists unique points $a_\theta, b_\theta \in \partial\mathcal{U}$ with $\eta_{a_\theta} = \theta$ and $\eta_{b_\theta} = -\theta$ the property that if we define $\mathcal{G}_\theta^{R_0} := \{z \in \mathcal{U} : \nabla \tilde{u}(z) \cdot R_0^{-1}\theta > 0\}$ we have

$$\left| \mathcal{U} \cap H\left(\frac{a_\theta + b_\theta}{2}, R_0\left(\frac{b_\theta - a_\theta}{|b_\theta - a_\theta|}\right)\right) \setminus \mathcal{G}_\theta^{R_0} \right| \leq c\beta^{\frac{1}{24}}. \quad (48)$$

Proof of Lemma 3. Without loss of generality assume Ω is centered on 0, i.e. $\int_\Omega z = 0$. Since $\partial\mathcal{U}$ is smooth and $\mathcal{U}$ is convex there exists a set $\Xi \subset S^1$ with $H^1(S^1 \setminus \Xi) = 0$ with the following property,

$$\exists \text{ unique } a_\varphi \in \partial\mathcal{U} \text{ with } \eta_{a_\varphi} = \varphi \text{ and a unique } b_\varphi \in \partial\mathcal{U} \text{ with } \eta_{b_\varphi} = -\varphi \text{ for all } \varphi \in \Xi. \quad (49)$$

Pick $\theta \in \Xi \cap \Gamma$ and let $\varphi := RR_0^{-1}\theta$ so note that $\varphi = \theta$ or $\varphi = -\theta$ depending on whether $R_0 = R$ or $R_0 = R^{-1}$.

Note since Ω is convex $\Omega \subset \overline{H(a_\varphi, \varphi)}$ we also know that $b_\varphi \in H(a_\varphi, \varphi)$ (since otherwise given that $\partial\Omega$ is smooth it would not be possible that $\eta_{b_\varphi} = -\varphi$), hence defining $\tau_\theta = \frac{b_\varphi - a_\varphi}{|b_\varphi - a_\varphi|}$ we have $\tau_\varphi \cdot \varphi > 0$.

Let $\tilde{m} = R(\nabla \tilde{u})$, it is easy to see that

$$\Pi_\varphi := \{z \in \mathcal{U} \setminus \Omega : \tilde{m}(z) \cdot \varphi > 0\} = \{z \in \mathcal{U} \setminus \Omega : \nabla u(z) \cdot R^{-1}\varphi > 0\} \quad (50)$$

forms a connected set whose boundary is contained in $\partial\mathcal{U}$ and $\partial\Omega$ and in two lines parallel to φ , see figure 2, also note the endpoints of $\partial\mathcal{U} \cap \overline{\Pi_\varphi}$ are given by a_φ and b_φ .

Now by Lemma 2, (41) function $\tilde{u}$ satisfies (6) and (7). Since either $\varphi = \theta \in \Gamma$ or $\varphi = -\theta \in \Gamma$ so we can apply Lemma 1, to $\tilde{m}$ and thus there exists function $w_\varphi : \mathcal{U} \rightarrow \mathbb{R}$ such that

$$\int_{\mathcal{U}} |\nabla w_\varphi - R(\Lambda_\varphi(\tilde{m}))| \leq c\beta^{\frac{1}{8}}. \quad (51)$$

By the Co-area formula and Chebyshev's inequality there exists a set $H \subset [0, 1/10]$ such that $H^1([0, 1/10] \setminus H) \leq c\beta^{\frac{1}{24}}$ where

$$\int_{\tilde{u}^{-1}(t)} |\nabla w_\varphi - R(\Lambda_\varphi(\tilde{m}))| dH^1 \leq c\beta^{\frac{1}{12}} \text{ for all } t \in H. \quad (52)$$

Pick $s_0 \in [1/10 - c\beta^{\frac{1}{24}}, 1/10] \cap H$. Recall $\tau_\varphi = \frac{b_\varphi - a_\varphi}{|b_\varphi - a_\varphi|}$ and

$$\mathcal{W}_\varphi := \overline{\mathcal{U}} \cap H\left(\frac{a_\varphi + b_\varphi}{2}, R\tau_\varphi\right). \quad (53)$$

We claim that

$$\partial\mathcal{U} \cap \overline{\Pi_\varphi} = \partial\mathcal{U} \cap \overline{\mathcal{W}_\varphi}. \quad (54)$$


Since the endpoints of $\partial\mathcal{U} \cap \overline{\Pi_\varphi}$ are the same as the endpoints of $\partial\mathcal{U} \cap \overline{\mathcal{W}_\varphi}$ it is sufficient to show $H^1(\partial\mathcal{U} \cap \overline{\Pi_\varphi} \cap \overline{\mathcal{W}_\varphi}) > 0$. Let

$$\Lambda = \sup \left\{ \lambda > 0 : \left(\frac{a_\varphi + b_\varphi}{2} + \lambda R\tau_\varphi + \langle \tau_\varphi \rangle \right) \cap \partial\mathcal{U} \neq \emptyset \right\}$$

then let c_φ be the point given by $\left(\frac{a_\varphi + b_\varphi}{2} + \Lambda R\tau_\varphi + \langle \tau_\varphi \rangle \right) \cap \partial\mathcal{U}$, since $\partial\mathcal{U}$ is smooth $\eta_{c_\varphi} = R^{-1}\tau_\varphi$, so $\nabla u(c_\varphi) = R^{-1}\tau_\varphi$ and thus $\nabla u(c_\varphi) \cdot R^{-1}\varphi = R^{-1}\tau_\varphi \cdot R^{-1}\varphi = \tau_\varphi \cdot \varphi > 0$. As this inequality is strict, in a neighborhood of c_φ the same inequality will be satisfied. Thus we have $H^1(\partial\mathcal{U} \cap \overline{\Pi_\varphi} \cap \overline{\mathcal{W}_\varphi}) > 0$ and so we have established (54).

By the construction of Π_φ and $\mathcal{W}_\varphi$ by (54) we have

$$H^1(\partial\Omega_{s_0} \cap \overline{\Pi_\varphi} \Delta \overline{\mathcal{W}_\varphi}) \leq c\beta^{\frac{1}{24}}. \quad (55)$$



1

FIGURE 2.

There must exist $\psi \in (0, 2\beta^{\frac{1}{24}})$ such that defining $Q := \begin{pmatrix} \cos \psi & -\sin \psi \\ \sin \psi & \cos \psi \end{pmatrix}$ we have

$$|R\varphi \cdot Q\tau_\varphi| > \beta^{\frac{1}{24}}. \quad (56)$$

Let $\zeta_\varphi := \frac{a_\varphi + b_\varphi}{2} + \mathcal{C}_2 \beta^{\frac{1}{24}} R\tau_\varphi$. From the construction it is clear that we can choose constant $\mathcal{C}_2$ large enough so that

$$\text{Card} \left(\partial\Omega_{s_0} \cap H \left(\frac{a_\varphi + b_\varphi}{2}, R\tau_\varphi \right) \cap \{\zeta_\varphi + \langle Q\tau_\varphi \rangle\} \right) = 2.$$

Now for every $t > 0$ for which $\partial\Omega_{s_0} \cap \{\zeta_\varphi + tR\tau_\varphi + \langle Q\tau_\varphi \rangle\} \neq \emptyset$ let ϱ_t^1, ϱ_t^2 be the points defined by $\{\varrho_t^1, \varrho_t^2\} = \partial\Omega_{s_0} \cap \{\zeta_\varphi + tR\tau_\varphi + \langle Q\tau_\varphi \rangle\}$ and $\varrho_t^2 \cdot \varphi \geq \varrho_t^1 \cdot \varphi$. By (55) we can assume constant $\mathcal{C}_2$ was chosen large enough so that $\varrho_t^1, \varrho_t^2 \in \Pi_\varphi$. Let Σ_t be the connected component

of $\partial\Omega_{s_0} \setminus \{\varrho_t^1, \varrho_t^2\}$ that lies inside Π_φ . Thus

$$\begin{aligned} |(w_\varphi(\varrho_t^2) - w_\varphi(\varrho_t^1)) - (\varrho_t^2 - \varrho_t^1) \cdot R\varphi| &= \left| \int_{\Sigma_t} \nabla w_\varphi(z) \cdot t_z dH^1 z - \int_{\Sigma_t} R\varphi \cdot t_z dH^1 z \right| \\ &= \left| \int_{\Sigma_t} (\nabla w_\varphi(z) - R\varphi) \cdot t_z dH^1 z \right| \\ &\stackrel{(52)}{\leq} c\beta^{\frac{1}{12}}. \end{aligned} \quad (57)$$

Let

$$e_t = \int_{[\varrho_t^1, \varrho_t^2]} |\nabla w_\varphi - R(\Lambda_\varphi(\tilde{m}))|, \quad (58)$$

so by the fundamental theorem of Calculus $\left| (w_\varphi(\varrho_t^2) - w_\varphi(\varrho_t^1)) - \int_{[\varrho_t^1, \varrho_t^2]} R(\Lambda_\varphi(\tilde{m})) \cdot Q\tau_\varphi \right| \leq e_t$. Thus in combination with (57) we have

$$\left| (\varrho_t^2 - \varrho_t^1) \cdot R\varphi - \int_{[\varrho_t^1, \varrho_t^2]} R(\Lambda_\varphi(\tilde{m})) \cdot Q\tau_\varphi \right| \leq e_t + c\beta^{\frac{1}{12}}. \quad (59)$$

Given the definition of Λ_φ (see (17)) and of $\mathcal{G}_\theta^{R_0}$ (see the statement of Lemma 3) so

$$R(\Lambda_\varphi(\tilde{m}(x))) = R\varphi \Leftrightarrow \tilde{m}(x) \cdot \varphi > 0 \Leftrightarrow \nabla \tilde{u}(x) \cdot R^{-1}\varphi > 0 \Leftrightarrow \nabla \tilde{u}(x) \cdot R_0^{-1}\theta > 0 \Leftrightarrow x \in \mathcal{G}_\theta^{R_0}.$$

In exactly the same way $\Lambda_\varphi(\tilde{m}(x)) = 0 \Leftrightarrow x \notin \mathcal{G}_\theta^{R_0}$. Hence

$$\int_{[\varrho_t^1, \varrho_t^2]} \Lambda_\varphi(\tilde{m}(x)) dH^1 x = \varphi H^1 \left([\varrho_t^1, \varrho_t^2] \cap \mathcal{G}_\theta^{R_0} \right)$$

which from (59)

$$\left| (\varrho_t^2 - \varrho_t^1) \cdot R\varphi - Q\tau_\varphi \cdot R\varphi H^1 \left([\varrho_t^1, \varrho_t^2] \cap \mathcal{G}_\theta^{R_0} \right) \right| \leq e_t + c\beta^{\frac{1}{12}}$$

since (recall (56)) we chose Q so that $|R\varphi \cdot Q\tau_\varphi| > \beta^{\frac{1}{24}}$ and since $\frac{\varrho_t^2 - \varrho_t^1}{|\varrho_t^2 - \varrho_t^1|} = Q\tau_\varphi$ so

$$\left| |\varrho_t^2 - \varrho_t^1| - H^1 \left([\varrho_t^1, \varrho_t^2] \cap \mathcal{G}_\theta^{R_0} \right) \right| \leq c\beta^{-\frac{1}{24}} e_t + c\beta^{\frac{1}{24}}.$$

To simplify notation let $\vartheta = H^1 (P_{\langle R(Q\tau_\varphi) \rangle} (\Omega_{s_0} \cap H(\zeta_\varphi, R(Q\tau_\varphi))))$

$$H^1 \left([\varrho_t^1, \varrho_t^2] \cap \mathcal{G}_\theta^{R_0} \right) \geq |\varrho_t^1 - \varrho_t^2| - c\beta^{-\frac{1}{24}} e_t - c\beta^{\frac{1}{24}} \text{ for any } t \in [0, \vartheta]. \quad (60)$$

So

$$\begin{aligned} \left| \Omega_{s_0} \cap H(\zeta_\varphi, R(Q\tau_\varphi)) \cap \mathcal{G}_\theta^{R_0} \right| &= \int_{[0, \vartheta]} H^1 \left([\varrho_t^1, \varrho_t^2] \cap \mathcal{G}_\theta^{R_0} \right) dt \\ &\stackrel{(60)}{\geq} \int_{[0, \vartheta]} |\varrho_t^1 - \varrho_t^2| - c\beta^{-\frac{1}{24}} e_t - c\beta^{\frac{1}{24}} \\ &\stackrel{(58)}{\geq} |\Omega_{s_0} \cap H(\zeta_\varphi, R(Q\tau_\varphi))| - c\beta^{\frac{1}{24}} \\ &\quad - c\beta^{-\frac{1}{24}} \int_{\mathcal{U}} |\nabla w_\varphi - R(\Lambda_\varphi(\tilde{m}))| \\ &\stackrel{(51)}{\geq} |\Omega_{s_0} \cap H(\zeta_\varphi, R(Q\tau_\varphi))| - c\beta^{\frac{1}{24}}. \end{aligned} \quad (61)$$

Note $|\mathcal{U} \setminus \Omega_{s_0}| \leq c\beta^{\frac{1}{24}}$ and by definition of $\mathcal{W}_\varphi$ (see (53)) $|\mathcal{W}_\varphi \setminus H(\zeta_\varphi, R(Q\tau_\varphi))| \leq c\beta^{\frac{1}{24}}$ this gives $|\mathcal{W}_\varphi \setminus \mathcal{G}_\theta^{R_0}| \leq c\beta^{\frac{1}{24}}$. Now if $R_0 = R$ and so $\varphi = \theta$, it is imediate that $\tau_\varphi = \frac{b_\theta - a_\theta}{|b_\theta - a_\theta|}$ and so (again recalling definition (53)) (48) follows. On the other hand if $R_0 = R^{-1}$ then $\varphi = -\theta$ and

so $a_\varphi = b_\theta$, $b_\varphi = a_\theta$, which implies $\tau_\varphi = -\frac{b_\theta - a_\theta}{|b_\theta - a_\theta|}$ so $R\tau_\varphi = R\left(-\frac{b_\theta - a_\theta}{|b_\theta - a_\theta|}\right) = R^{-1}\left(\frac{b_\theta - a_\theta}{|b_\theta - a_\theta|}\right) = R_0\left(\frac{b_\theta - a_\theta}{|b_\theta - a_\theta|}\right)$ hence (again recalling definition (53)), (48) also follows in this case. $\square$

Lemma 4. *Let Ω be a convex body centered on 0 with $\text{diam}(\Omega) = 2$. Let $u : W^{2,2}(\Omega) \rightarrow \mathbb{R}$ be a function satisfying (6) and (7) and in addition u satisfies $u = 0$ on $\partial\Omega$ and $\nabla u(z) \cdot \eta_z = 1$ on $\partial\Omega$ in the sense of trace where η_z is the inward pointing unit normal to $\partial\Omega$ at z . Let $a, b \in \Omega$ be such that $\text{diam}(\Omega) = |a - b|$ we will show there exists constant $\mathcal{C}_3 > 1$ and $r_0 \in (\mathcal{C}_3^{-1}\beta^{\frac{1}{256}}, \mathcal{C}_3\beta^{\frac{1}{256}})$ such that*

$$u(x) \geq 1 - \mathcal{C}_3\beta^{\frac{1}{256}} \text{ for any } x \in \partial B_{r_0}\left(\frac{a+b}{2}\right). \quad (62)$$

Proof of Lemma 4. Let $\mathcal{U}$ be the convex set and $\tilde{u}$ be the function constructed in Lemma 3. It is easy to see we can chose $\tilde{a}, \tilde{b} \in \mathcal{U}$ such that $\frac{\tilde{a}-\tilde{b}}{|\tilde{a}-\tilde{b}|} = \frac{a-b}{|a-b|}$ and $|\tilde{a} - \tilde{b}| = \text{diam}(\mathcal{U})$. Without loss of generality assume $\frac{\tilde{a}-\tilde{b}}{|\tilde{a}-\tilde{b}|} = e_2$.

Step 1. Let $P : [0, H^1(\partial\mathcal{U})) \rightarrow \partial\mathcal{U}$ be a ‘clockwise’ parameterisation of $\mathcal{U}$ by arclength with $P(0) = \tilde{a}$. Let $\sigma_1 = P(H^1(\partial\mathcal{U}) - \beta^{\frac{1}{256}})$ and $\sigma_2 = P(\beta^{\frac{1}{256}})$, see figure 3. The points σ_1, σ_2 satisfy the following properties, firstly

$$\eta_{\sigma_i} \cdot (-e_2) \geq 1 - c\beta^{\frac{1}{128}} \text{ for } i = 1, 2. \quad (63)$$

Secondly

$$|\sigma_1 - \sigma_2| \leq 2\beta^{\frac{1}{256}}. \quad (64)$$

Thirdly

$$\sigma_1 \cdot (-e_1) \geq \frac{\beta^{\frac{1}{256}}}{2} \text{ and } \sigma_2 \cdot e_1 \geq \frac{\beta^{\frac{1}{256}}}{2}. \quad (65)$$

Proof of Step 1. Recall $\mathcal{U} = \Omega_{\frac{1}{10}}(\Omega)$, so for any $x \in \partial\mathcal{U}$ let $z_x \in \partial\Omega$ be such that $d(x, \Omega) = |x - z_x|$, note that we can inscribe a ball $B_{\frac{1}{10}}(z_x) \subset \mathcal{U}$ with $x \in \partial B_{\frac{1}{10}}(z_x) \cap \partial\mathcal{U}$ and $B_{\frac{1}{10}}(z_x) \cap \partial\mathcal{U} = \emptyset$. Thus the curvature of $\partial\mathcal{U}$ is bounded above by 10 and so $\|\ddot{P}\|_{L^\infty(\partial\mathcal{U})} \leq 10$. Since $\dot{P}(\tilde{a}) = e_1$, $|\dot{P}(\tilde{a}) - \dot{P}(\beta^{\frac{1}{256}})| \leq 10\beta^{\frac{1}{256}}$ and as $\eta_{\sigma_2} = R^{-1}(\dot{P}(\beta^{\frac{1}{256}}))$ this proves (63) for $i = 2$. The proof for $i = 1$ follows the same way.

Inequality (64) follows instantly since σ_1 and σ_2 is connected by a path of length less than $2\beta^{\frac{1}{256}}$. Now from for any $t \in [0, \beta^{\frac{1}{256}}]$ we have

$$|e_1 - \dot{P}(t)| = |\dot{P}(0) - \dot{P}(t)| \leq \int_0^t |\ddot{P}(s)| ds \leq 10\beta^{\frac{1}{256}}.$$

Thus $\sigma_2 \cdot e_1 = (\sigma_2 - \tilde{a}) \cdot e_1 = \int_0^{\beta^{\frac{1}{256}}} \dot{P}(s) \cdot e_1 ds \geq (1 - 10\beta^{\frac{1}{256}})\beta^{\frac{1}{256}}$. Arguing in exactly the same way $(-e_1) \cdot \sigma_1 \geq (1 - 10\beta^{\frac{1}{256}})\beta^{\frac{1}{256}}$ which establishes (65).

Step 2. We will show there exists positive constant $\mathcal{C}_4$ and $x_0 \in N_{\mathcal{C}_4\beta^{\frac{1}{256}}}([\tilde{a}, \tilde{b}]) \cap \mathcal{U}$ such that for some $\psi_0 \in B_{\mathcal{C}_4\beta^{\frac{1}{256}}}(e_2)$ the following inequality holds

$$\left| X\left(x_0, \psi_0, \mathcal{C}_4\beta^{\frac{1}{256}}\right) \cap \mathcal{U} \setminus \left\{x : |\nabla \tilde{u}(x) \cdot e_1| < \mathcal{C}_4\beta^{\frac{1}{256}}\right\} \right| \leq \mathcal{C}_4\beta^{\frac{1}{24}}. \quad (66)$$

Proof of Step 2. Without loss of generality we can assume σ_1 and σ_2 are chosen so that $\eta_{\sigma_1} \in \Xi$ and $\eta_{\sigma_2} \in \Xi$ (recall the definition of Ξ , see (49)). We also know $\eta_{\tilde{a}} = -e_2$ and $\eta_{\tilde{b}} = e_2$. Let $\omega_1 \in \partial\mathcal{U}$ be the unique point for which $-\eta_{\omega_1} = \eta_{\sigma_1}$ and let $\omega_2 \in \partial\mathcal{U}$ be the unique point for which $-\eta_{\omega_2} = \eta_{\sigma_2}$, see figure 3.

Define

$$\Pi_2 := H\left(\frac{\sigma_2 + \omega_2}{2}, R\left(\frac{\omega_2 - \sigma_2}{|\omega_2 - \sigma_2|}\right)\right) \cap H\left(\frac{\sigma_1 + \omega_1}{2}, R^{-1}\left(\frac{\omega_1 - \sigma_1}{|\omega_1 - \sigma_1|}\right)\right) \quad (67)$$

and

$$\Pi_1 := H\left(\frac{\sigma_2 + \omega_2}{2}, R^{-1}\left(\frac{\omega_2 - \sigma_2}{|\omega_2 - \sigma_2|}\right)\right) \cap H\left(\frac{\sigma_1 + \omega_1}{2}, R\left(\frac{\omega_1 - \sigma_1}{|\omega_1 - \sigma_1|}\right)\right) \quad (68)$$

and let $\Pi = \Pi_1 \cup \Pi_2$ and let $x_0 := \overline{\Pi_1} \cap \overline{\Pi_2}$, see figure 3.




FIGURE 3.

Let us define $l_x^\theta := x + \mathbb{R}_+\theta$ for any $x \in \mathbb{R}^2$, $\theta \in S^1$. First we will show $(x_0 + \mathbb{R}e_2) \subset \Pi$ however this inclusion is relatively easy to see because firstly

$$e_2 \cdot R\left(\frac{\omega_1 - \sigma_1}{|\omega_1 - \sigma_1|}\right) = e_1 \cdot \left(\frac{\omega_1 - \sigma_1}{|\omega_1 - \sigma_1|}\right) \geq \frac{\beta^{\frac{1}{256}}}{4}$$

thus $l_0^{e_2} \subset H\left(0, R\left(\frac{\omega_1 - \sigma_1}{|\omega_1 - \sigma_1|}\right)\right)$. And secondly as $x_0 \in \partial H\left(\frac{\sigma_1 + \omega_1}{2}, R\left(\frac{\omega_1 - \sigma_1}{|\omega_1 - \sigma_1|}\right)\right)$

$$l_{x_0}^{e_2} \subset H\left(x_0, R\left(\frac{\omega_1 - \sigma_1}{|\omega_1 - \sigma_1|}\right)\right) = H\left(\frac{\sigma_1 + \omega_1}{2}, R\left(\frac{\omega_1 - \sigma_1}{|\omega_1 - \sigma_1|}\right)\right).$$

In exactly the same way $l_{x_0}^{e_2} \subset H\left(\frac{\sigma_2 + \omega_2}{2}, R^{-1}\left(\frac{\omega_2 - \sigma_2}{|\omega_2 - \sigma_2|}\right)\right)$. Hence $l_{x_0}^{e_2} \subset \Pi_1$. Arguing in the same manner we have $l_{x_0}^{e_2} \subset \Pi_2$ and thus we have established the claim.

Let $\gamma = l_{x_0}^{e_2} \cap \partial\mathcal{U}$, by construction we have that γ lies in the component of $\partial\mathcal{U}$ between σ_1 and σ_2 and hence we know $d(\gamma, l_0^{e_2}) \leq c\beta^{\frac{1}{256}}$ and so it follows $x_0 \in N_{c\beta^{\frac{1}{256}}}\left(\left[\tilde{a}, \tilde{b}\right]\right) \cap \mathcal{U}$.

Since $\eta_{\tilde{a}} = -e_2$, $\eta_{\tilde{b}} = e_2$ and $\mathcal{U}$ is convex we know $\omega_2 \in H(0, -e_1)$ and for the same reasons $\omega_1 \in H(0, e_1)$ see figure 3. So $(\sigma_2 - \omega_2) \cdot e_1 \geq \sigma_2 \cdot e_1 \geq c\beta^{\frac{1}{256}}$ and for exactly the same reason $(\sigma_1 - \omega_1) \cdot (-e_1) \geq \sigma_2 \cdot (-e_1) \geq c\beta^{\frac{1}{256}}$. Thus as $|\sigma_1 - \omega_1| \leq 2\text{diam}(\mathcal{U})$ and $|\sigma_2 - \omega_2| \leq 2\text{diam}(\mathcal{U})$ we have $\frac{\sigma_2 - \omega_2}{|\sigma_2 - \omega_2|} \cdot e_1 \geq c\beta^{\frac{1}{256}}$ and $\frac{\sigma_1 - \omega_1}{|\sigma_1 - \omega_1|} \cdot (-e_1) \geq c\beta^{\frac{1}{256}}$. Hence

$$\begin{aligned} \left(\frac{\sigma_1 - \omega_1}{|\sigma_1 - \omega_1|}\right) \cdot \left(\frac{\sigma_2 - \omega_2}{|\sigma_2 - \omega_2|}\right) &= \left(\frac{\sigma_1 - \omega_1}{|\sigma_1 - \omega_1|} \cdot e_1\right) \left(\frac{\sigma_2 - \omega_2}{|\sigma_2 - \omega_2|} \cdot e_1\right) \\ &\quad + \left(\frac{\sigma_1 - \omega_1}{|\sigma_1 - \omega_1|} \cdot e_2\right) \left(\frac{\sigma_2 - \omega_2}{|\sigma_2 - \omega_2|} \cdot e_2\right) \\ &\leq -c\beta^{\frac{1}{128}} + 1. \end{aligned}$$

In other words the angle between $\frac{\sigma_1 - \omega_1}{|\sigma_1 - \omega_1|}$ and $\frac{\sigma_2 - \omega_2}{|\sigma_2 - \omega_2|}$ is greater than $\mathcal{C}_4\beta^{\frac{1}{256}}$ for some positive constant $\mathcal{C}_4$. Thus there exists $\psi_0 \in B_{c\beta^{\frac{1}{256}}}(e_2)$ such that $X\left(x_0, \psi_0, \mathcal{C}_4\beta^{\frac{1}{256}}\right) \subset \Pi$. By Lemma 3 we know that

$$\left|\mathcal{U} \cap H\left(\frac{\sigma_2 + \omega_2}{2}, R^{-1}\left(\frac{\omega_2 - \sigma_2}{|\omega_2 - \sigma_2|}\right)\right) \setminus \mathcal{G}_{\eta_{\sigma_2}}^{R^{-1}}\right| \leq c\beta^{\frac{1}{24}}$$

and

$$\left|\mathcal{U} \cap H\left(\frac{\sigma_1 + \omega_1}{2}, R\left(\frac{\omega_1 - \sigma_1}{|\omega_1 - \sigma_1|}\right)\right) \setminus \mathcal{G}_{\eta_{\sigma_1}}^R\right| \leq c\beta^{\frac{1}{24}}.$$

Thus (recalling the definition of Π_1 , (68))

$$\left|\Pi_1 \cap \mathcal{U} \setminus \mathcal{G}_{\eta_{\sigma_2}}^{R^{-1}} \cap \mathcal{G}_{\eta_{\sigma_1}}^R\right| \leq c\beta^{\frac{1}{24}}. \quad (69)$$

In exactly the same way we have (recall (67))

$$\left|\Pi_2 \cap \mathcal{U} \setminus \mathcal{G}_{\eta_{\sigma_1}}^{R^{-1}} \cap \mathcal{G}_{\eta_{\sigma_2}}^R\right| \leq c\beta^{\frac{1}{24}}. \quad (70)$$

Now for any $x \in \mathcal{G}_{\eta_{\sigma_2}}^{R^{-1}} \cap \mathcal{G}_{\eta_{\sigma_1}}^R$ we have $\nabla \tilde{u}(x) \cdot R\eta_{\sigma_2} \geq 0$ and $\nabla \tilde{u}(x) \cdot R^{-1}\eta_{\sigma_1} \geq 0$. Since from (63) $\eta_{\sigma_i} \in X^+\left(0, -e_2, c\beta^{\frac{1}{256}}\right)$ for $i = 1, 2$ we know $R\eta_{\sigma_2} \in X^+\left(0, e_1, c\beta^{\frac{1}{256}}\right)$ and $R^{-1}\eta_{\sigma_1} \in X^+\left(0, -e_1, \beta^{\frac{1}{256}}\right)$, from this it is easy to see (assuming we chose $\mathcal{C}_4$ large enough) $|\nabla \tilde{u}(x) \cdot e_1| \leq \mathcal{C}_4\beta^{\frac{1}{256}}$. And in the same way for any $x \in \mathcal{G}_{\eta_{\sigma_1}}^{R^{-1}} \cap \mathcal{G}_{\eta_{\sigma_2}}^R$ we also have $|\nabla \tilde{u}(x) \cdot e_1| \leq \mathcal{C}_4\beta^{\frac{1}{256}}$.

$$\begin{aligned} &\left|X\left(x_0, \psi_0, \mathcal{C}_4\beta^{\frac{1}{256}}\right) \cap \mathcal{U} \setminus \left\{x : |\nabla u(x) \cdot e_1| < \mathcal{C}_4\beta^{\frac{1}{256}}\right\}\right| \\ &\leq c\left|\Pi_1 \cap \mathcal{U} \setminus \mathcal{G}_{\eta_{\sigma_1}}^R \cap \mathcal{G}_{\eta_{\sigma_2}}^{R^{-1}}\right| + c\left|\Pi_2 \cap \mathcal{U} \setminus \mathcal{G}_{\eta_{\sigma_2}}^R \cap \mathcal{G}_{\eta_{\sigma_1}}^{R^{-1}}\right| \\ &\leq \mathcal{C}_4\beta^{\frac{1}{24}} \end{aligned}$$

which establishes (66).

Step 3. There exists positive constant $\mathcal{C}_5$ such that for some $v_1 \in \{e_2, -e_2\}$ we have

$$\left| X \left(x_0, \psi_0, \mathcal{C}_4 \beta^{\frac{1}{256}} \right) \cap \mathcal{U} \cap H \left(\mathcal{C}_5 \beta^{\frac{1}{256}} v_1, v_1 \right) \setminus \mathbb{V}_{-v_1} \right| \leq \mathcal{C}_5 \beta^{\frac{1}{24}}. \quad (71)$$

where

$$\mathbb{V}_{-v_1} := \left\{ x \in \mathcal{U} : \nabla u(x) \in N_{\mathcal{C}_5 \beta^{\frac{1}{256}}}(-v_1) \right\}. \quad (72)$$

Proof of Step 3. Let $\varpi_0 = l_0^{-e_1} \cap \partial \mathcal{U}$. Note since $\mathcal{U}$ is convex $\eta_{\varpi_0} \cdot e_1 > 0$. We claim

$$\eta_{\varpi_0} \cdot e_1 > \frac{1}{10}. \quad (73)$$

Suppose this were not the case, then $\eta_{\varpi_0} \cdot e_1 \leq \frac{1}{10}$. Since $\mathcal{U}$ is convex and $\text{diam}(\mathcal{U}) < 2$ we know $\mathcal{U} \subset \overline{H(\varpi_0, \eta_{\varpi_0})} \subset H(-2e_1, \eta_{\varpi_0})$ which implies $(\tilde{b} + 2e_1) \cdot \eta_{\varpi_0} > 0$ and thus

$$\tilde{b} \cdot e_2 \sqrt{\frac{99}{100}} \geq \left((\tilde{b} + 2e_1) \cdot e_2 \right) (\eta_{\varpi_0} \cdot e_2) > - \left((\tilde{b} + 2e_1) \cdot e_1 \right) (\eta_{\varpi_0} \cdot e_1) = -2\eta_{\varpi_0} \cdot e_1 \geq -\frac{1}{5}$$

however as $|\tilde{a} - \tilde{b}| = \text{diam}(\mathcal{U}) = \frac{22}{10}$, $\frac{\tilde{a} + \tilde{b}}{2} = 0$ and $\frac{\tilde{a} - \tilde{b}}{|\tilde{a} - \tilde{b}|} = e_2$ this is a contradiction.

Let $\varpi_1 \in \partial \mathcal{U}$ a point for which $\eta_{\varpi_1} = -\eta_{\varpi_0}$. Now let $l \in \left(\frac{\varpi_1 - \varpi_0}{|\varpi_1 - \varpi_0|} \right)^\perp \cap S^1$ be such that

$$H^1 \left([a, b] \cap H \left(\frac{\varpi_1 + \varpi_0}{2}, l \right) \right) \geq \frac{|a - b|}{2}. \quad (74)$$

Chose $S \in \{R^{-1}, R\}$ so that $S \left(\frac{\varpi_1 - \varpi_0}{|\varpi_1 - \varpi_0|} \right) = l$, by Lemma 3 we have

$$\left| \mathcal{U} \cap H \left(\frac{\varpi_1 + \varpi_0}{2}, l \right) \setminus \mathcal{G}_{\eta_{\varpi_0}}^S \right| \leq c\beta^{\frac{1}{24}}. \quad (75)$$

From (7) and (66) we know

$$\left| X \left(x_0, \psi_0, \mathcal{C}_4 \beta^{\frac{1}{256}} \right) \cap \mathcal{U} \setminus \{x : \nabla \tilde{u}(x) \in N_{100^{-1}}(\{e_2, -e_2\})\} \right| \leq c\beta^{\frac{1}{24}}. \quad (76)$$

Since so $|S^{-1}\eta_{\varpi_0} \cdot e_2| \stackrel{(73)}{>} 10^{-1}$ there exists some fixed vector $v_0 \in \{e_2, -e_2\}$ such that if $x \in \mathcal{G}_{\eta_{\varpi_0}}^S \cap \{x : \nabla \tilde{u}(x) \in N_{100^{-1}}(\{e_2, -e_2\})\}$ then $\nabla \tilde{u}(x) \in B_{100^{-1}}(v_0)$ and so using (75) and (76)

$$\left| X \left(x_0, \psi_0, \mathcal{C}_4 \beta^{\frac{1}{256}} \right) \cap \mathcal{U} \cap H \left(\frac{\varpi_1 + \varpi_0}{2}, l \right) \setminus \{x : \nabla \tilde{u}(x) \in B_{100^{-1}}(v_0)\} \right| \leq c\beta^{\frac{1}{24}}. \quad (77)$$

Now for any $w \in H(0, v_0)$ we have the elementary inequality $|w - v_0| \leq 4d(w, S^1) + |w \cdot e_1|$, so using (7), (66) and (77) we have (assuming constant $\mathcal{C}_5$ is large enough, recall definition (72))

$$\left| X \left(x_0, \psi_0, \mathcal{C}_4 \beta^{\frac{1}{256}} \right) \cap \mathcal{U} \cap H \left(\frac{\varpi_1 + \varpi_0}{2}, l \right) \setminus \mathbb{V}_{v_0} \right| \leq c\beta^{\frac{1}{24}}. \quad (78)$$

Recall $\varpi_0 = l_0^{-e_1} \cap \partial \mathcal{U}$ so as $B_{\frac{1}{10}}(0) \subset \mathcal{U}$ we have that $\varpi_0 \cdot (-e_1) > \frac{1}{10}$. Now either $\varpi_1 \cdot e_1 > -\frac{1}{20}$ and so $\left| \frac{\varpi_0 - \varpi_1}{|\varpi_0 - \varpi_1|} \cdot e_1 \right| \geq \frac{1}{40}$ and so $|l \cdot e_2| \geq \frac{1}{40}$. Thus by the fact that $\psi_0 \in B_{\mathcal{C}_4 \beta^{\frac{1}{256}}}(e_2)$ and that inequality (74) implies $0 \in \overline{H(\frac{\varpi_1 + \varpi_0}{2}, l)}$ there exists $v_1 \in \{e_2, -e_2\}$ such that for some constant $\mathcal{C}_5$ we have

$$X \left(x_0, \psi_0, \mathcal{C}_4 \beta^{\frac{1}{256}} \right) \cap \mathcal{U} \cap H \left(\mathcal{C}_5 \beta^{\frac{1}{256}} v_1, v_1 \right) \subset H \left(\frac{\varpi_1 + \varpi_0}{2}, l \right). \quad (79)$$

On the other hand if $\varpi_1 \cdot e_1 \leq -\frac{1}{20}$ then by (74) we know $H \left(\frac{\varpi_1 + \varpi_0}{2}, l \right)$ is the larger part of $\mathcal{U}$ and indeed $X \left(x_0, \psi_0, \mathcal{C}_4 \beta^{\frac{1}{256}} \right) \cap \mathcal{U}$ will be fully contained in $H \left(\frac{\varpi_1 + \varpi_0}{2}, l \right)$ and this also implies (79).

Putting (79) together with (78) gives

$$\left| X \left(x_0, \psi_0, \mathcal{C}_4 \beta^{\frac{1}{256}} \right) \cap \mathcal{U} \cap H \left(\mathcal{C}_5 \beta^{\frac{1}{256}} v_1, v_1 \right) \setminus \mathbb{V}_{v_0} \right| \leq c \beta^{\frac{1}{24}}.$$

Let $x \in \mathcal{U} \setminus \overline{\Omega} \cap X \left(x_0, \psi_0, \mathcal{C}_4 \beta^{\frac{1}{256}} \right) \cap H \left(\mathcal{C}_5 \beta^{\frac{1}{256}} v_1, v_1 \right)$ so as $\tilde{u}(x) = d(x, \partial \mathcal{U})$ and since $\psi_0 \in B_{\mathcal{C}_4 \beta^{\frac{1}{256}}}(e_2)$ so $\nabla \tilde{u}(x) \in N_{\mathcal{C}_5 \beta^{\frac{1}{256}}}(-v_1)$ thus we must have $v_0 = -v_1$, this gives (71).

Step 4. We will show there exists a positive constant $\mathcal{C}_6$ such that

$$(l_x^{-\theta} \cup l_x^{\theta}) \setminus B_{\mathcal{C}_6 \beta^{\frac{1}{128}}}(x) \subset X(x_0, \psi_0, \mathcal{C}_4 \beta^{\frac{1}{256}}) \text{ for all } x \in B_{\frac{1}{128}}(x_0), \theta \in S^1 \cap B_{\beta^{\frac{1}{128}}}(\psi_0). \quad (80)$$

Proof of Step 4. Without loss of generality we assume $x_0 = 0$, $\psi_0 = e_2$ and $\mathcal{C}_4 = 1$. To begin with to take point $x = \beta^{\frac{1}{128}} e_1$, we will show later the general case follows from this. See figure 4.

Let $\theta = \left(\frac{\sin \beta^{\frac{1}{128}}}{\cos \beta^{\frac{1}{128}}} \right)$ and let $y = \partial X(0, e_2, \beta^{\frac{1}{256}}) \cap l_x^{\theta}$. We will get an upper bound on $|y|$. Let $z = y \cdot e_1 e_1$. We have two triangles to calculate with, triangle T_1 with corners on $0, x, y$ which is a subset of triangle T_2 with corners on $0, z, y$. Note that by applying the law of sines we have $|y|^{-1} \sin(\frac{\pi}{2} + \beta^{\frac{1}{128}}) = |x - y|^{-1} \sin(\frac{\pi}{2} - \beta^{\frac{1}{256}})$. Note that $T_3 = T_2 \setminus T_1$ is also a right angle triangle and since $|z| = \beta^{\frac{1}{128}} + |x - z|$ we have $|y| \cos(\frac{\pi}{2} - \beta^{\frac{1}{256}}) = \beta^{\frac{1}{128}} + |y - x| \cos(\frac{\pi}{2} - \beta^{\frac{1}{128}})$. Putting this together with the previous equation we have $|y| \sin \beta^{\frac{1}{256}} = \beta^{\frac{1}{128}} + |y| \frac{\cos \beta^{\frac{1}{256}}}{\cos \beta^{\frac{1}{128}}} \sin \beta^{\frac{1}{128}}$ which gives $|y| \left(\sin \beta^{\frac{1}{256}} - \frac{\cos \beta^{\frac{1}{256}}}{\cos \beta^{\frac{1}{128}}} \sin \beta^{\frac{1}{128}} \right) = \beta^{\frac{1}{128}}$. Now by taking the Taylor series approximating $\sin$ and $\cos$ we have $|y| \left(\beta^{\frac{1}{256}} + O \left(\beta^{\frac{1}{128}} \right) \right) = \beta^{\frac{1}{128}}$. Thus $|y| \sim \beta^{\frac{1}{256}}$ and thus the existence of constant $\mathcal{C}_6$ such that (80) holds follows instantly for the case $x = \beta^{\frac{1}{128}} e_1$.

In the general case where $x \neq \beta^{\frac{1}{128}} e_1$ suppose without loss of generality $x \cdot e_1 > 0$, define $\tilde{x} = (x + \langle \theta \rangle) \cap \langle e_1 \rangle$, since the angle between θ and e_1 is with $c \beta^{\frac{1}{256}}$ of $\frac{\pi}{2}$ it is easy to see $\tilde{x} \in B_{2\beta^{\frac{1}{128}}}(0)$ and of course $l_x^{\theta} \cap \partial X(0, e_2, \beta^{\frac{1}{256}}) = l_{\tilde{x}}^{\theta} \cap \partial X(0, e_2, \beta^{\frac{1}{256}})$ so the argument for the special case $x = \beta^{\frac{1}{128}} e_1$ can be applied to show the existence of constant $\mathcal{C}_6$ satisfying (80).

Step 5. We will establish (62).

Proof of Step 5. Let


$$h(z) := \mathbb{1}_{X(x_0, \psi_0, \mathcal{C}_4 \beta^{\frac{1}{256}}) \cap H(\mathcal{C}_5 \beta^{\frac{1}{256}} v_1, v_1) \cap \mathcal{U} \setminus \mathbb{V}_{-v_1}} \quad (81)$$

so we know $\int h \stackrel{(71)}{\leq} c \beta^{\frac{1}{24}}$. So by the Fubini's Theorem

$$\begin{aligned} & \int_{\mathcal{U}} \int_{\mathcal{U}} \left(h(z) + \beta^{-1} \left| 1 - |\nabla \tilde{u}(z)|^2 \right|^2 \right) |z - x|^{-1} dz dx \\ & \leq \int_{\mathcal{U}} \left(h(z) + \beta^{-1} \left| 1 - |\nabla \tilde{u}(z)|^2 \right|^2 \right) \left(\int |z - x|^{-1} dx \right) dz \\ & \leq c \int_{\mathcal{U}} \left(h(z) + \beta^{-1} \left| 1 - |\nabla \tilde{u}(z)|^2 \right|^2 \right) dz \\ & \stackrel{(7)}{\leq} c \beta^{\frac{1}{24}}. \end{aligned} \quad (82)$$

Let

$$G := \left\{ x \in B_{\beta^{\frac{1}{128}}}(x_0) : \int_{\mathcal{U}} \left(h(z) + \beta^{-1} \left| 1 - |\nabla \tilde{u}(z)|^2 \right|^2 \right) |z - x|^{-1} dz \leq c \beta^{\frac{1}{48}} \right\} \quad (83)$$



1

FIGURE 4.

so we know $\beta^{\frac{1}{48}} \left| B_{\beta^{\frac{1}{128}}}(x_0) \setminus G \right| \leq c\beta^{\frac{1}{24}}$, thus $\left| B_{\beta^{\frac{1}{128}}}(x_0) \setminus G \right| \leq c\beta^{\frac{1}{48}}$, assuming β is small enough $|G| \geq 2^{-1}\beta^{\frac{1}{64}}$. By Step 4, (80) for any $x \in B_{\beta^{\frac{1}{128}}}(x_0)$, $\theta \in B_{\beta^{\frac{1}{128}}}(\psi_0) \cap S^1$ we have $(l_x^{-\theta} \cup l_x^{\theta}) \setminus B_{C_6\beta^{\frac{1}{256}}}(x) \subset X(x_0, \psi_0, C_4\beta^{\frac{1}{256}})$.

Since $X(0, \psi_0, C_4\beta^{\frac{1}{256}}) = X(0, -\psi_0, C_4\beta^{\frac{1}{256}})$ we can assume without loss generality that $\psi_0 \cdot v_1 > 0$. Pick $x \in G$, by the Co-area formula we must be able to find $\theta_1 \in B_{\beta^{\frac{1}{128}}}(\psi_0) \cap S^1$ such that

$$\int_{(l_x^{-\theta_1} \cup l_x^{\theta_1}) \cap \mathcal{U}} h(z) + \beta^{-1} \left| 1 - |\nabla \tilde{u}(z)|^2 \right|^2 dH^1 z \leq c\beta^{\frac{1}{48}} / \beta^{-\frac{1}{128}} \leq c\beta^{\frac{1}{128}}. \quad (84)$$

So let d, e be the endpoints of the connected set $1d$ set $(l_x^{-\theta_1} \cup l_x^{\theta_1}) \cap \mathcal{U} \cap H(\mathcal{C}_5 \beta^{\frac{1}{256}} v_1, v_1)$ where we chose $d \in \partial H(\mathcal{C}_5 \beta^{\frac{1}{256}} v_1, v_1)$ and $e \in \partial \mathcal{U}$, note by Step 4 $[d, e] \setminus B_{\mathcal{C}_6 \beta^{\frac{1}{128}}}(x) \subset X(x_0, \psi_0, \mathcal{C}_4 \beta^{\frac{1}{256}}) \cap H(\mathcal{C}_5 \beta^{\frac{1}{256}} v_1, v_1)$. Now since for any $z \in [d, e] \setminus B_{\mathcal{C}_6 \beta^{\frac{1}{128}}}(x)$ with $h(z) = 0$ by (81) we have $z \in \mathbb{V}_{-v_1}$ so

$$H^1([d, e] \setminus \mathbb{V}_{-v_1}) \leq 2\mathcal{C}_6 \beta^{\frac{1}{128}} + H^1\left([d, e] \setminus (B_{\mathcal{C}_6 \beta^{\frac{1}{128}}}(x) \cup \mathbb{V}_{-v_1})\right) \stackrel{(84)}{\leq} c\beta^{\frac{1}{128}}. \quad (85)$$

Note also that if $z \in \mathbb{V}_{-v_1}$ so $\nabla \tilde{u}(z) \in B_{\mathcal{C}_5 \beta^{\frac{1}{256}}}(-v_1)$ and as $\theta_1 \in B_{\beta^{\frac{1}{128}}}(\psi_0) \subset B_{\mathcal{C}_4 \beta^{\frac{1}{256}}}(v_1)$ thus $\nabla \tilde{u}(z) \cdot (-\theta_1) \geq 1 - c\beta^{\frac{1}{128}}$ and by the fundamental theorem of Calculus

$$\begin{aligned} \tilde{u}(d) - \tilde{u}(e) &= \int_{[e, d]} \nabla u(z) \cdot (-\theta_1) dH^1 z \\ &\geq \int_{\mathbb{V}_{-v_1} \cap [e, d]} \nabla \tilde{u}(z) \cdot (-\theta_1) dH^1 z - \int_{[e, d] \setminus \mathbb{V}_{-v_1}} |\nabla \tilde{u}(z)| dH^1 z \\ &\stackrel{(85)}{\geq} \left(1 - c\beta^{\frac{1}{128}}\right) H^1(\mathbb{V}_{-v_1} \cap [e, d]) - H^1([e, d] \setminus \mathbb{V}_{-v_1}) \\ &\quad - c \int_{[e, d]} |1 - |\nabla \tilde{u}|^2| dH^1 \\ &\stackrel{(84)}{\geq} |d - e| (1 - c\beta^{\frac{1}{128}}). \end{aligned} \quad (86)$$

Since the curvature of $\partial \mathcal{U}$ is bounded above by 10 it is easy to see that

$$|e - \tilde{a}| \leq c\beta^{\frac{1}{256}}, \quad (87)$$

it is also easy to see $[e, \tilde{a}] \subset \mathcal{U} \setminus \Omega$ and $\tilde{u}$ is 1-Lipschitz on $\mathcal{U} \setminus \Omega$ so

$$|\tilde{u}(e) - \tilde{u}(\tilde{a})| \leq c\beta^{\frac{1}{256}}. \quad (88)$$

Thus we have

$$\begin{aligned} \tilde{u}(d) &= \tilde{u}(d) - \tilde{u}(\tilde{a}) \\ &\stackrel{(86), (87), (88)}{\geq} |d - \tilde{a}| - c\beta^{\frac{1}{256}} \\ &\geq |\tilde{a}| - c\beta^{\frac{1}{256}} = 2^{-1} \text{diam}(\mathcal{U}) - c\beta^{\frac{1}{256}}. \end{aligned} \quad (89)$$

Pick $r_0 \in [|d|, |d| + \beta^{\frac{1}{256}}]$ such that $\int_{\partial B_{r_0}(0)} |1 - |\nabla \tilde{u}(z)|^2| dH^1 z \leq c\beta^{-\frac{1}{256}} \beta$. Now fix $y \in \partial B_{r_0}(0)$, let $s = [d, e] \cap \partial B_{r_0}(0)$ and Γ_1 denote a connected component of $\partial B_{r_0}(0) \setminus \{s, y\}$. So we know $\int_{\Gamma_1 \cup [d, s]} |\nabla \tilde{u}(z)| dH^1 z \leq cH^1(\Gamma_1 \cup [d, s]) \leq c\beta^{\frac{1}{256}}$ so we can apply the fundamental theorem of Calculus we have that $|u(y) - u(d)| \leq c\beta^{\frac{1}{256}}$ and since y was an arbitrary point in $\partial B_{r_0}(0)$, using (89) this gives

$$\inf \{\tilde{u}(z) : z \in \partial B_{r_0}(0)\} \geq 2^{-1} \text{diam}(\mathcal{U}) - c\beta^{\frac{1}{256}}. \quad (90)$$

By definition (see (42)) $\tilde{u}(z) = u(z) + 10^{-1}$ for any $z \in \partial B_{r_0}(0)$. Since $\text{diam}(\mathcal{U}) = \frac{22}{10}$ putting this with (90) we have (62). $\square$

Proof of Theorem 2. Let $r_0 \in (\mathcal{C}_3^{-1} \beta^{\frac{1}{256}}, \mathcal{C}_3 \beta^{\frac{1}{256}})$ be a number we obtain from Lemma 4 that satisfies (62). By Fubini's Theorem we know $\int_{\Omega} \int_{\Omega} |1 - |\nabla u(z)|^2|^2 |z - y|^{-1} dz dy \leq \mathcal{C}_7 \beta^2$ for

some constant $\mathcal{C}_7 > 0$. Let

$$G_0 := \left\{ y \in \Omega : \int_{\Omega} \left| 1 - |\nabla u(z)|^2 \right|^2 |z - y|^{-1} dz \leq \beta \right\}. \quad (91)$$

Note that $|\Omega \setminus G_0| \leq \mathcal{C}_7 \beta$.

Let $a, b \in \overline{\Omega}$ be such that $|a - b| = \text{diam}(\Omega)$. Let $\vartheta = \frac{a+b}{2}$. Since $r_0 > \mathcal{C}_3^{-1} \beta^{\frac{1}{256}}$ we can pick $x_0 \in B_{\beta^{\frac{1}{4}}}(\vartheta) \cap G_0 \subset B_{r_0}(\vartheta)$. So by the Co-area formula there exists $\Psi \subset S^1$ such that $H^1(S^1 \setminus \Psi) \leq \sqrt{\beta}$ and

$$\int_{l_{x_0}^{\theta}} \left| 1 - |\nabla u|^2 \right|^2 dH^1 z \leq c\sqrt{\beta} \text{ for each } \theta \in \Psi. \quad (92)$$

For any $\theta \in S^1$ define $P(\theta) := l_{x_0}^{\theta} \cap \partial\Omega$, we will show

$$|P(\theta) - x_0| \geq 1 - c\beta^{\frac{1}{256}} \text{ for any } \theta \in \Psi. \quad (93)$$

To see this we argue as follows, for each $z \in [x, P(\theta)]$ let $\theta_z \in S^1$ be such that $|\nabla u(z) - \theta_z| = d(\nabla u(z), S^1)$. Note $d(\nabla u(z), S^1) \leq c||\nabla u(z)| - 1|$

$$\begin{aligned} \int_{[x_0, P(\theta)]} |\nabla u(z) - \theta_z| dH^1 z &\leq c \left(\int_{[x_0, P(\theta)]} \left| |\nabla u(z)|^2 - 1 \right|^2 dH^1 z \right)^{\frac{1}{2}} \\ &\stackrel{(92)}{\leq} c\beta^{\frac{1}{4}}. \end{aligned} \quad (94)$$

So

$$\begin{aligned} u(x_0) &= u(x_0) - u(P(\theta)) \\ &= \int_{[x_0, P(\theta)]} \nabla u(z) \cdot (-\theta) dH^1 z \\ &\stackrel{(94)}{\leq} \int_{[x_0, P(\theta)]} \theta_z \cdot (-\theta) dH^1 z + c\beta^{\frac{1}{4}} \\ &\leq |x_0 - P(\theta)| + c\beta^{\frac{1}{4}}. \end{aligned} \quad (95)$$

Let $y_{\theta} := [x_0, P(\theta)] \cap \partial B_{r_0}(\vartheta)$. In exactly the same way we have

$$|u(y_{\theta}) - u(x_0)| \leq c\beta^{\frac{1}{256}}. \quad (96)$$

So

$$u(x_0) \geq u(y_{\theta}) - |u(y_{\theta}) - u(x_0)| \stackrel{(96)}{\geq} u(y_{\theta}) - c\beta^{\frac{1}{256}} \stackrel{(62)}{\geq} 1 - c\beta^{\frac{1}{256}} \quad (97)$$

this together with (95) establishes (93).

Let $N = \left\lceil 2^{-1} \beta^{-\frac{1}{2}} \right\rceil$, we can divide S^1 into N disjoint pieces of equal length, denote them $I_1, I_2, \dots, I_N$. Formally; $\bigcup_{k=1}^N I_k = S^1$ and $H^1(I_k) = \frac{2\pi}{N}$ for each $k = 1, 2, \dots, N$. We can pick $\theta_k \in I_k \cap \Psi$ for each $k = 1, 2, \dots, N$.

Let

$$h = \min \{ |P(\theta_k) - x_0| : k \in \{1, 2, \dots, N\} \}. \quad (98)$$

We define Π to be the convex hull of the points $x_0 + h\theta_1, x_0 + h\theta_2, \dots, x_0 + h\theta_N$. Now by the construction of Π , for any $y \in \partial\Pi$ we can find $k \in \{1, 2, \dots, N\}$ such that $|y - (x_0 + h\theta_k)| \leq c\sqrt{\beta}$ and thus $|y - x_0| \geq h - c\sqrt{\beta}$ and so

$$B_{h-c\sqrt{\beta}}(x_0) \subset \Pi. \quad (99)$$

Note that by using (93) we know $h > 1 - c\beta^{\frac{1}{256}}$ and since $|x_0 - \vartheta| \leq \beta^{\frac{1}{4}}$ (recalling also that Ω is convex and so $\Pi \subset \Omega$) there exists positive constant $\mathcal{C}_8$ such that

$$B_{1-\mathcal{C}_8\beta^{\frac{1}{256}}}(\vartheta) \subset \Omega. \quad (100)$$

We claim

$$\Omega \subset B_{1+2\mathcal{C}_8\beta^{\frac{1}{256}}}(\vartheta). \quad (101)$$

Suppose not, so there exists $y \in \partial\Omega$ such that $|y - \vartheta| \geq 1 + 2\mathcal{C}_8\beta^{\frac{1}{256}}$. By inequality (100) we know $-\frac{y-\vartheta}{|y-\vartheta|} \left(1 - \mathcal{C}_8\beta^{\frac{1}{256}}\right) + \vartheta \in \Omega$ and as by convexity of Ω , $\left[y, \vartheta - \frac{y-\vartheta}{|y-\vartheta|} \left(1 - \mathcal{C}_8\beta^{\frac{1}{256}}\right)\right] \subset \Omega$ thus

$$H^1 \left(\left[y, \vartheta - \frac{y-\vartheta}{|y-\vartheta|} \left(1 - \mathcal{C}_8\beta^{\frac{1}{256}}\right) \right] \right) \geq 2 + \mathcal{C}_8\beta^{\frac{1}{256}}$$

which contradicts the fact $\text{diam}(\Omega) = 2$ hence (101) is established. Since the center of mass of Ω is 0, i.e. $\int_{\Omega} dx = 0$, by (100), (101) we have that $|\vartheta| \leq c\beta^{\frac{1}{256}}$. Now $|x_0 - P(\theta)| \leq |P(\theta)| + |x_0| \stackrel{(101)}{\leq} 1 + c\beta^{\frac{1}{256}}$ so putting this together with (97) we have

$$u(x_0) - u(P(\theta)) = u(x_0) \geq |x_0 - P(\theta)| - c\beta^{\frac{1}{256}}. \quad (102)$$

Thus

$$\begin{aligned} \int_{[x_0, P(\theta)]} |\nabla u(z) + \theta|^2 dH^1 z &= \int_{[x_0, P(\theta)]} (|\nabla u(z)|^2 + 2\nabla u(z) \cdot \theta + 1) dH^1 z \\ &\stackrel{(92)}{\leq} 2(1 + c\beta^{\frac{1}{4}}) |x_0 - P(\theta)| + 2(u(P(\theta)) - u(x_0)) \\ &\stackrel{(102)}{\leq} c\beta^{\frac{1}{256}} \text{ for any } \theta \in \Psi. \end{aligned} \quad (103)$$

Now using the elementary fact that $\left| \nabla u(z) + \frac{z-x_0}{|z-x_0|} \right|^2 \leq \left| |\nabla u(z)|^2 - 1 \right|^2 + c$, since $x_0 \in G_0$ we have

$$\begin{aligned} \int_{\theta \in S^1 \setminus \Psi} \int_{I_{x_0}^{\theta}} \left| \nabla u(z) + \frac{z-x_0}{|z-x_0|} \right|^2 dH^1 z dH^1 \theta \\ \leq cH^1(S^1 \setminus \Psi) + \int_{\theta \in S^1} \int_{I_{x_0}^{\theta}} \left| |\nabla u(z)|^2 - 1 \right|^2 dH^1 z dH^1 \theta \\ \stackrel{(92)}{\leq} c\sqrt{\beta}. \end{aligned} \quad (104)$$

And thus

$$\begin{aligned} \int_{\Omega} \left| \nabla u(z) + \frac{z-x_0}{|z-x_0|} \right|^2 dz &\leq \int_{\Omega} \left| \nabla u(z) + \frac{z-x_0}{|z-x_0|} \right|^2 |z-x_0|^{-1} dz \\ &\leq \int_{\theta \in S^1} \int_{I_{x_0}^{\theta}} \left| \nabla u(z) + \frac{z-x_0}{|z-x_0|} \right|^2 dH^1 z dH^1 \theta \\ &\stackrel{(104), (103)}{\leq} c\beta^{\frac{1}{256}}. \end{aligned}$$

By Holder's inequality this gives

$$\left(\int_{\Omega} \left| \nabla u(z) + \frac{z-x_0}{|z-x_0|} \right|^2 dz \right)^{\frac{1}{2}} \leq c\beta^{\frac{1}{512}}. \quad (105)$$

Now for any $z \in \Omega \setminus B_{\beta^{\frac{1}{512}}}(0)$, since $|x_0| \leq c\beta^{\frac{1}{256}}$

$$\begin{aligned} \left| \frac{z}{|z|} - \frac{z-x_0}{|z-x_0|} \right| &= \left| \frac{z|z-x_0| - (z-x_0)|z|}{|z||z-x_0|} \right| \\ &= \left| \frac{z(|z-x_0|-|z|) + x_0|z|}{|z||z-x_0|} \right| \\ &\leq \left| \frac{|z-x_0|-|z|}{|z-x_0|} \right| + \frac{|x_0|}{|z-x_0|} \\ &\leq c\beta^{\frac{1}{512}}. \end{aligned} \tag{106}$$

So

$$\left(\int_{\Omega} \left| \frac{z}{|z|} - \frac{z-x_0}{|z-x_0|} \right|^2 dz \right)^{\frac{1}{2}} \leq c\beta^{\frac{1}{256}} + \left(\int_{\Omega \setminus B_{\beta^{\frac{1}{512}}}} \left| \frac{z}{|z|} - \frac{z-x_0}{|z-x_0|} \right|^2 dz \right)^{\frac{1}{2}} \stackrel{(106)}{\leq} c\beta^{\frac{1}{512}}.$$

Putting this together with (105) we have (8). $\square$

4. CONSTRUCTION OF UPPER BOUND FOR NEARLY CIRCULAR DOMAINS

In this section we will show that given a convex domain Ω with C^2 boundary with curvature bounded above by $\epsilon^{-\frac{1}{2}}$ and that satisfies $|B_1(0) \triangle \Omega| \leq \beta$ we will construct a function u with $I_{\epsilon}(u) \leq \beta^{\frac{3}{16}}$, this is the contents of Proposition 1 below. The proof of Corollary 1 will follow easily from this.

Proposition 1. *Let Ω be a convex body with C^2 boundary and with curvature bounded above by $\epsilon^{-\frac{1}{2}}$ and $|\Omega \triangle B_1(0)| \leq \beta$. Let $\epsilon \in (0, \frac{\beta^{\frac{1}{2}}}{4}]$, there exists a function C^∞ function $\xi : \Omega \rightarrow \mathbb{R}$ which satisfies $\nabla u(z) \cdot \eta_z = 1$ (where η_z is the inward pointing unit normal to $\partial\Omega$ at z) and for which*

$$\int_{\Omega} \epsilon^{-1} \left| 1 - |\nabla \xi|^2 \right|^2 + \epsilon |\nabla^2 \xi|^2 dz \leq c\beta^{\frac{3}{32}}. \tag{107}$$

4.1. Proof of Proposition 1. We begin with a preliminary lemma

Lemma 5. *Let $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a continuous function. Let ρ denote the standard convolution kernel, i.e. $\int \rho = 1$ and $\text{Spt} \rho \subset B_1(0)$ and define $\rho_h(z) := h^{-2} \rho(h^{-1}z)$.*

*Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be an affine function with $\eta = \nabla f$, let $g(x) = f * \rho_{\phi(x)}(x)$ then*

$$g(x) = f(x) \text{ for all } x \in \mathbb{R}^n. \tag{108}$$

Proof. So

$$\begin{aligned} g(x) &= \int f(x-y)(\phi(x))^{-2} \rho(\phi(x)^{-1}y) dy \\ &= \int (f(x) + \eta \cdot (x-y))(\phi(x))^{-2} \rho(\phi(x)^{-1}y) dy \\ &= f(x) + \int \eta \cdot (x-y)(\phi(x))^{-2} \rho(\phi(x)^{-1}y) dy \\ &= f(x) + \int_{[-\phi(x)+x \cdot \eta, \phi(x)+x \cdot \eta]} \int_{\lambda \eta + \eta^\perp} (x \cdot \eta - \lambda)(\phi(x))^{-2} \rho(\phi(x)^{-1}z) dH^{n-2} z d\lambda \\ &= f(x) + \int_{[-\phi(x), 0]} \int_{(x \cdot \eta + \lambda) \eta + \eta^\perp} -\lambda(\phi(x))^{-2} \rho(\phi(x)^{-1}z) dH^{n-2} z d\lambda \\ &\quad + \int_{[0, \phi(x)]} \int_{(x \cdot \eta + \lambda) \eta + \eta^\perp} -\lambda(\phi(x))^{-2} \rho(\phi(x)^{-1}z) dH^{n-2} z d\lambda \end{aligned} \tag{109}$$

Since for any $\lambda \in [0, \phi(x)]$ we have

$$\int_{(x \cdot \eta - \lambda)\eta + \eta^\perp} \lambda(\phi(x))^{-2} \rho(\phi(x)^{-1}z) dH^{n-2}z = \int_{(x \cdot \eta + \lambda)\eta + \eta^\perp} \lambda(\phi(x))^{-2} \rho(\phi(x)^{-1}z) dH^{n-2}z$$

thus the last two term of (109) cancel and so $g(x) = f(x)$. This completes the proof of the lemma. $\square$

Lemma 6. *Suppose Ω is a convex and $|\Omega \triangle B_1| = \beta$. Let $a_\theta = \partial\Omega \cap l_0^\theta$ we have*

$$||a_\theta| - 1| \leq c\sqrt{\beta} \text{ and so } \partial\Omega \subset N_{c\sqrt{\beta}}(\partial B_1(0)). \quad (110)$$

In addition for any θ there exists constant c such that

$$|\eta_{a_\theta} + \theta| \leq c\beta^{\frac{1}{4}}. \quad (111)$$

Proof of Lemma.

Step 1. We will show $B_{\frac{1}{2}}(0) \subset \Omega$.

Proof of Step 1. Suppose not, so we can pick $x \in \partial\Omega \cap B_{\frac{1}{2}}(0)$. Let η_x be an inward pointing unit normal to $\partial\Omega$ at x , by convexity of Ω we have $\Omega \subset \overline{H(x, \eta_x)}$ and so $B_1(0) \cap H(x, -\eta_x) \cap \Omega = \emptyset$ which implies $|B_1(0) \setminus \Omega| \geq |B_1(0) \cap H(x, -\eta_x)| > \frac{1}{8}$ which contradicts that $|\Omega \triangle B_1| \leq \beta$.

Step 2. $a_\theta \in B_{1+c\sqrt{\beta}}(0)$.

Proof of Step 2. Suppose not. Since Ω is convex we have $\text{conv}(\{a_\theta\} \cup B_{\frac{1}{2}}(0)) \subset \Omega$ and

$$|\text{conv}(\{a_\theta\} \cup B_{\frac{1}{2}}(0)) \setminus B_1(0)| > c\beta,$$

thus we have $|\Omega \setminus B_1(0)| > c\beta$ which contradicts the fact that $|\Omega \triangle B_1(0)| = \beta$.

Step 3. We will show $a_\theta \notin B_{1-c\sqrt{\beta}}(0)$.

Proof of Step 3. Suppose $a_\theta \in B_{1-c\sqrt{\beta}}(0)$ this implies $|B_1(0) \setminus H(a_\theta, \eta_{a_\theta})| \geq c\beta$ and $\Omega \subset H(a_\theta, \eta_{a_\theta})$ so $|B_1(0) \setminus \Omega| \geq c\beta$ which gives a contradiction.

Proof of Lemma completed. Suppose (111) is false, since $|a_\theta - \theta| \leq c\sqrt{\beta}$ we have

$$|B_1(0) \setminus H(a_\theta, \eta_{a_\theta})| \geq c\beta^{\frac{3}{4}},$$

as before this implies $|B_1(0) \setminus \Omega| > c\beta^{\frac{3}{4}}$ which is a contradiction. $\square$

Lemma 7. *Let Ω be convex and define $u(x) := d(z, \partial\Omega)$ for any $z \in \Omega$ then function u is concave*

Proof of Lemma. Let $a, b \in \Omega$. Since Ω is convex $\text{conv}(B_{u(a)}(a) \cup B_{u(b)}(b)) \subset \Omega$. Now suppose there exists $\lambda \in (0, 1)$ such that

$$u(\lambda a + (1 - \lambda)b) < \lambda u(a) + (1 - \lambda)u(b)$$

then as this implies $B_{u(\lambda a + (1 - \lambda)b)}(\lambda a + (1 - \lambda)b) \subset \text{int}(\text{conv}(B_{u(a)}(a) \cup B_{u(b)}(b)))$ we must be able to find $x \in \partial\Omega$ with $x \in \partial\Omega \cap \text{conv}(B_{u(a)}(a) \cup B_{u(b)}(b))$ which is a contradiction. $\square$

Lemma 8. *Let $\epsilon > 0$, suppose Ω is a convex body with C^2 boundary and with curvature bounded above by $\epsilon^{-\frac{1}{2}}$. We will construct a function $\psi : \Omega \cap N_\epsilon(\partial\Omega) \rightarrow \mathbb{R}$ with the following properties*

$$\int_{\Omega \cap N_\epsilon(\partial\Omega)} |1 - |\nabla\psi|^2|^2 \leq c\epsilon^2, \quad (112)$$

$$\int_{\Omega \cap N_\epsilon(\partial\Omega)} |\nabla^2\psi|^2 \leq c, \quad (113)$$

$$\psi(z) = [u * \rho_\epsilon](z) \text{ for any } z \in \Omega \setminus N_\epsilon(\partial\Omega) \quad (114)$$

and

$$\nabla\psi(z) = \eta_z \text{ for each } z \in \partial\Omega. \quad (115)$$

Proof. Let $w : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a smooth monotonic function with the following properties

$$w(z) = \begin{cases} z & \text{for } z \in [0, \frac{\epsilon}{3}) \\ \epsilon & \text{for } z \geq \epsilon \end{cases} \quad (116)$$

and $\sup |\ddot{w}| \leq c\epsilon^{-1}$.

Let $u(x) = d(x, \partial\Omega)$. For any $x \in \Omega \cap N_\epsilon(\partial\Omega)$ define $\phi(x) = w(u(x))$. Let ρ be the standard convolution kernel, i.e. as defined in Lemma 5. We will convolve the function u with convolution kernel $\rho_{\phi(x)}(z) := \rho\left(\frac{z}{\phi(x)}\right) / (\phi(x))^2$. Since the convolution kernel varies with x , when we differentiate $u * \rho_{\phi(x)}$, the derivative will involve a term with the derivative of $\rho_{\phi(x)}$. For this reason we need to calculate various partial derivatives of $\rho_{\phi(x)}$.

For each $x \in \Omega \cap N_\epsilon(\partial\Omega)$ let $b_x \in \partial\Omega$ be defined by $|x - b_x| = u(x)$. Note that since the curvature of $\partial\Omega$ is bounded above by $\epsilon^{-\frac{1}{2}}$, for any $x \in \Omega \cap N_\epsilon(\partial\Omega)$ we have that there is one unique $b_x \in \partial\Omega$ such that $|x - b_x| = u(x)$. We define $\varsigma_x = \frac{x - b_x}{|x - b_x|}$, let $R = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and define $\omega_x = R\varsigma_x$.

Note $\varsigma_x = \eta_{b_x}$, i.e. the inward pointing unit normal to $\partial\Omega$ at b_x . Note also that for all small enough h , $b_x = b_{x+h\varsigma_x}$ so $u(x+h\varsigma_x) = h + u(x)$. Thus

$$\begin{aligned} \phi_{,\varsigma_x}(x) &= \lim_{h \rightarrow 0} \frac{\phi(x+h\varsigma_x) - \phi(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{w(u(x)+h) - w(u(x))}{h} \\ &= \dot{w}(u(x)). \end{aligned}$$

Note also that since $|\nabla u(x)| = 1$ and $u_{,\varsigma_x}(x) = \lim_{h \rightarrow 0} \frac{u(x+h\varsigma_x) - u(x)}{h} = 1$ so

$$u_{,\omega_x}(x) = \lim_{h \rightarrow 0} \frac{u(x+h\omega_x) - u(x)}{h} = 0.$$

Thus

$$\phi_{,\omega_x}(x) = \dot{w}(u(x))u_{,\omega_x}(x) = 0. \quad (117)$$

So

$$\begin{aligned} \frac{\partial}{\partial \varsigma_x}(\rho_{\phi(x)}(z)) &= \frac{\partial}{\partial \varsigma_x} \left(\rho\left(\frac{z}{\phi(x)}\right) (\phi(x))^{-2} \right) \\ &= -\nabla \rho\left(\frac{z}{\phi(x)}\right) \cdot z \frac{\phi_{,\varsigma_x}(x)}{(\phi(x))^4} - 2\rho\left(\frac{z}{\phi(x)}\right) \frac{\phi_{,\varsigma_x}(x)}{(\phi(x))^3} \end{aligned} \quad (118)$$

and

$$\frac{\partial}{\partial \omega_x}(\rho_{\phi(x)}(z)) = 0. \quad (119)$$

Define $\psi(x) := \int u(x-z)\rho_{\phi(x)}(z)dz$. Now

$$\begin{aligned} \psi_{,\varsigma_x}(x) &\stackrel{(118)}{=} \int u_{,\varsigma_x}(x-z)\rho_{\phi(x)}(z)dz \\ &\quad - \int u(x-z) \left(\nabla \rho\left(\frac{z}{\phi(x)}\right) \cdot z \frac{\phi_{,\varsigma_x}(x)}{(\phi(x))^4} + 2\rho\left(\frac{z}{\phi(x)}\right) \frac{\phi_{,\varsigma_x}(x)}{(\phi(x))^3} \right) dz \end{aligned} \quad (120)$$

In the same way it is easy to see $\psi_{,\omega_x}(x) = 0$ and so

$$\psi_{,\varsigma_x \omega_x}(x) = 0. \quad (121)$$

We also know that $u_{,\varsigma_x \varsigma_x}(x) = 0$ so

$$\begin{aligned} \psi_{,\varsigma_x \varsigma_x}(x) &= \int u(x-z) \partial_{\varsigma_x} \left(\sum_{k=1}^2 -\rho_{,k} \left(\frac{z}{\phi(x)} \right) \frac{z_k \phi_{,\varsigma_x}(x)}{(\phi(x))^4} - 2\rho \left(\frac{z}{\phi(x)} \right) \frac{\phi_{,\varsigma_x}(x)}{(\phi(x))^3} \right) dz \\ &= \int u(x-z) \left(\sum_{k,l=1}^2 \rho_{,kl} \left(\frac{z}{\phi(x)} \right) \frac{(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^6} z_k z_l - \sum_{k=1}^2 \rho_{,k} \left(\frac{z}{\phi(x)} \right) z_k \partial_{\varsigma_x} \left(\frac{\phi_{,\varsigma_x}(x)}{(\phi(x))^4} \right) \right. \\ &\quad \left. + 2 \sum_{m=1}^2 \rho_{,m} \left(\frac{z}{\phi(x)} \right) z_m \frac{(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^4} - 2\rho \left(\frac{z}{\phi(x)} \right) \partial_{\varsigma_x} \left(\frac{\phi_{,\varsigma_x}(x)}{(\phi(x))^3} \right) \right) dz \end{aligned}$$

Note

$$\partial_{\varsigma_x} \left(\frac{\phi_{,\varsigma_x}(x)}{(\phi(x))^3} \right) = \frac{-3(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^4} + \frac{\phi_{,\varsigma_x \varsigma_x}(x)}{(\phi(x))^3}$$

and

$$\partial_{\varsigma_x} \left(\frac{\phi_{,\varsigma_x}(x)}{(\phi(x))^4} \right) = \frac{-4(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^5} + \frac{\phi_{,\varsigma_x \varsigma_x}(x)}{(\phi(x))^4}. \quad (122)$$

So

$$\begin{aligned} \psi_{,\varsigma_x \varsigma_x}(x) &= \int u(x-z) \left(\left(\nabla^2 \rho \left(\frac{z}{\phi(x)} \right) : z \otimes z \right) \frac{(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^6} \right. \\ &\quad \left. + \left(-\frac{\phi_{,\varsigma_x \varsigma_x}(x)}{(\phi(x))^4} + \frac{4(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^5} + \frac{2(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^4} \right) \nabla \rho \left(\frac{z}{\phi(x)} \right) \cdot z \right. \\ &\quad \left. + \left(\frac{6(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^4} - \frac{2\phi_{,\varsigma_x \varsigma_x}(x)}{(\phi(x))^3} \right) \rho \left(\frac{z}{\phi(x)} \right) \right) dz. \end{aligned} \quad (123)$$

Step 1. For any $x \in N_\epsilon(\partial\Omega) \cap \Omega$ we have

$$\sup \{ |\nabla u(z) - \varsigma_x| : z \in B_{4u(x)}(x) \cap \Omega \} \leq c\epsilon^{-\frac{1}{2}} u(x). \quad (124)$$

Proof of Step 1. Since $\partial\Omega$ has curvature less than $\epsilon^{-\frac{1}{2}}$ for any $x_1, x_2 \in \partial\Omega$, $[x_1, x_1 + \epsilon^{\frac{1}{2}}\eta_{x_1}] \cap [x_2, x_2 + \epsilon^{\frac{1}{2}}\eta_{x_2}] = \emptyset$. So for any $x_1, x_2 \in B_{4u(x)}(x) \cap \partial\Omega$, $|\eta_{x_1} - \eta_{x_2}| \leq \epsilon^{-\frac{1}{2}} H^1(B_{4u(x)}(x) \cap \partial\Omega)$. Note as $\Omega \cap B_{4u(x)}(x)$ is convex and $\partial\Omega \cap B_{4u(x)}(x) \subset \partial(\Omega \cap B_{4u(x)}(x))$ so $H^1(\partial\Omega \cap B_{4u(x)}(x)) \leq cu(x)$. Hence $|\eta_{x_1} - \eta_{x_2}| \leq c\epsilon^{-\frac{1}{2}} u(x) \leq c\sqrt{\epsilon}$ so it is clear that

$$B_{4u(x)}(x) \cap \Omega \subset \bigcup_{z \in \partial\Omega \cap B_{4u(x)}(x)} [z, z + \sqrt{\epsilon}\eta_z]. \quad (125)$$

For any $z \in B_{4u(x)}(x) \cap \Omega$ we have $\nabla u(z) = \frac{z-b_z}{|z-b_z|} = \eta_{b_z}$ where b_z is such that $|z-b_z| = d(z, \partial\Omega)$. So for any $z_1, z_2 \in B_{4u(x)}(x) \cap \Omega$ by (125) we have that $b_{z_1}, b_{z_2} \in \partial\Omega \cap B_{4u(x)}(x)$, so $|\nabla u(z_1) - \nabla u(z_2)| = |\eta_{b_{z_1}} - \eta_{b_{z_2}}| \leq c\epsilon^{-\frac{1}{2}} u(x)$.

Step 2. For any $x \in N_\epsilon(\partial\Omega) \cap \Omega$ we have

$$||\nabla \psi(x)| - 1| \leq c\sqrt{\epsilon}. \quad (126)$$

And

$$\lim_{y \rightarrow z} \nabla \psi(y) = \eta_z. \quad (127)$$

Proof of Step 2. From (120) we have

$$\begin{aligned}
& |\psi_{\varsigma_x}(x) - 1| \\
& \leq \left| \overbrace{\int (u_{\varsigma_x}(x-z) - 1) \rho\left(\frac{z}{\phi(x)}\right) (\phi(x))^{-2} dz}^B \right| \\
& \quad + \left| \overbrace{\int \frac{-u(x-z)\phi_{\varsigma_x}(x)}{(\phi(x))^3} \left(\nabla \rho\left(\frac{z}{\phi(x)}\right) \cdot \frac{z}{\phi(x)} + 2\rho\left(\frac{z}{\phi(x)}\right) \right) dz}^C \right|. \quad (128)
\end{aligned}$$

Now for any $z \in \text{Spt}\rho_{\phi(x)}$ we have that $\nabla u(x-z) = u_{\varsigma_x}(x-z)\varsigma_x + u_{\omega_x}(x-z)\omega_x$ now since $\text{Spt}\rho_{\phi(x)} \subset B_{\phi(x)}(0) \subset B_{u(x)}(0)$ so for any $z \in \text{Spt}\rho_{\phi(x)}$ by (124) from Step 1 we have $|\nabla u(x-z) - \varsigma_x| \leq c\epsilon^{-\frac{1}{2}}u(x)$ and thus

$$|u_{\varsigma_x}(x-z) - 1| \leq c\epsilon^{-\frac{1}{2}}u(x) \quad (129)$$

so (noting $u(x) \leq c\phi(x)$ for any $x \in N_\epsilon(\partial\Omega) \cap \Omega$)

$$B \leq cu(x)\epsilon^{-\frac{1}{2}} < c\phi(x)\epsilon^{-\frac{1}{2}}. \quad (130)$$

Also defining $w = \oint_{B_{\phi(x)}(x)} \nabla u$

$$|w - \varsigma_x| = \left| \oint_{B_{\phi(x)}(x)} (\nabla u(z) - \varsigma_x) dz \right| \stackrel{(124)}{\leq} c\epsilon^{-\frac{1}{2}}\phi(x). \quad (131)$$

So by Poincare inequality there exists affine function l_w with $\nabla l_w = w$

$$\begin{aligned}
\oint_{B_{\phi(x)}(x)} |u(z) - l_w(z)| dz & \leq c\phi(x) \oint_{B_{\phi(x)}(x)} |\nabla u(z) - w| dz \\
& \leq c\phi(x) \left(\oint_{B_{\phi(x)}(x)} |\nabla u(z) - \varsigma_x| dz + c|w - \varsigma_x| \right) \\
& \stackrel{(124), (131)}{\leq} c\epsilon^{-\frac{1}{2}}(\phi(x))^2. \quad (132)
\end{aligned}$$

Now using (131), again for the appropriate choice of affine function l_{ς_x} with $\nabla l_{\varsigma_x} = \varsigma_x$ we have by Poincare's inequality

$$\oint_{B_{\phi(x)}(x)} |l_{\varsigma_x}(z) - l_w(z)| dz \leq c\phi(x) \oint_{B_{\phi(x)}(x)} |w - \varsigma_x| dz \stackrel{(131)}{\leq} c\epsilon^{-\frac{1}{2}}(\phi(x))^2$$

with (132) gives

$$\oint_{B_{\phi(x)}(x)} |l_{\varsigma_x}(z) - u(z)| dz \leq c\epsilon^{-\frac{1}{2}}(\phi(x))^2. \quad (133)$$

Let g be defined by $g(x) = l_{\varsigma_x} * \rho_{\phi(x)}(x)$, note by Lemma 5 we have $\nabla g(z) = \varsigma_x$ for any $z \in \mathbb{R}^2$ and hence $g_{\varsigma_x}(x) = 1$ and as

$$\begin{aligned}
g_{\varsigma_x}(x) &= \int \rho\left(\frac{z}{\phi(x)}\right) (\phi(x))^{-2} dz \\
&\quad - \int \frac{l_{\varsigma_x}(x-z)}{(\phi(x))^3} \phi_{\varsigma_x}(x) \left(\nabla \rho\left(\frac{z}{\phi(x)}\right) \cdot z (\phi(x))^{-1} + 2\rho\left(\frac{z}{\phi(x)}\right) \right) dz \\
&= 1 - \int \frac{l_{\varsigma_x}(x-z)}{(\phi(x))^3} \phi_{\varsigma_x}(x) \left(\nabla \rho\left(\frac{z}{\phi(x)}\right) \cdot z (\phi(x))^{-1} + 2\rho\left(\frac{z}{\phi(x)}\right) \right) dz.
\end{aligned}$$

Thus

$$0 = \int \frac{l_{\varsigma_x}(x-z)}{(\phi(x))^3} \phi_{,\varsigma_x}(x) \left(\nabla \rho \left(\frac{z}{\phi(x)} \right) \cdot z (\phi(x))^{-1} + 2\rho \left(\frac{z}{\phi(x)} \right) \right) dz \quad (134)$$

So

$$\begin{aligned} C &\leq \int \frac{|l_{\varsigma_x}(x-z) - u(x-z)|}{(\phi(x))^3} \left| \phi_{,\varsigma_x}(x) \left(\nabla \rho \left(\frac{z}{\phi(x)} \right) \cdot z (\phi(x))^{-1} + 2\rho \left(\frac{z}{\phi(x)} \right) \right) \right| dz \\ &\leq c(\phi(x))^{-3} \int_{B_{\phi(x)}(x)} |l_{\varsigma_x}(z) - u(z)| dz \\ (133) \quad &\leq c\epsilon^{-\frac{1}{2}} \phi(x). \end{aligned} \quad (135)$$

Since $x \in N_\epsilon(\partial\Omega) \cap \Omega$ we know $\phi(x) \leq c\epsilon$ applying (135) and (130) to (128) gives

$$|\psi_{,\varsigma_x}(x) - 1| \leq c\epsilon^{-\frac{1}{2}} \phi(x) \leq c\sqrt{\epsilon}. \quad (136)$$

As $\psi_{,\omega_x}(x) = 0$, so $|\nabla\psi(x) - \varsigma_x| \leq c\sqrt{\epsilon}$ and (126) follows easily. Also for (136) we know $|\nabla\psi(x) - \eta_{b_x}| \leq c\epsilon^{-\frac{1}{2}}u(x)$ and (127) follows. This completes the proof of Step 2.

Step 3. For any $x \in N_\epsilon(\partial\Omega) \cap \Omega$ we have $|\nabla^2\psi(x)| \leq c\epsilon^{-\frac{1}{2}}$.

Proof Step 3. From Step 2 (133) we know the existence of an affine function l_{ς_x} with $\nabla l_{\varsigma_x} = \varsigma_x$ with $\int_{B_{\phi(x)}(x)} |u - l_{\varsigma_x}| dz \leq c\epsilon^{\frac{1}{2}} \phi(x)^2$.

Let $g(x) := l_{\varsigma_x} * \rho_{\phi(x)}(x)$ so by Lemma 5 we know $g_{,\varsigma_x\varsigma_x}(x) = 0$. By following through the same calculation that gave (123) we have

$$\begin{aligned} 0 &= \int l_{\varsigma_x}(x-z) \left(\left(\nabla^2 \rho \left(\frac{z}{\phi(x)} \right) : z \otimes z \right) \frac{(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^6} \right. \\ &\quad + \left(-\frac{\phi_{,\varsigma_x\varsigma_x}(x)}{(\phi(x))^4} + \frac{4(\phi_{,\varsigma_x}(x))^4}{(\phi(x))^5} + \frac{2(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^4} \right) \nabla \rho \left(\frac{z}{\phi(x)} \right) \cdot z \\ &\quad \left. + \left(\frac{6(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^4} - \frac{2\phi_{,\varsigma_x\varsigma_x}(x)}{(\phi(x))^3} \right) \rho \left(\frac{z}{\phi(x)} \right) \right) dz. \end{aligned} \quad (137)$$

So applying this to (123)

$$\begin{aligned} &|\psi_{,\varsigma_x\varsigma_x}(x)| \\ &\leq \int |u(x-z) - l_{\varsigma_x}(x-z)| \left| \left(\nabla^2 \rho \left(\frac{z}{\phi(x)} \right) : z \otimes z \right) \frac{(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^6} \right. \\ &\quad + \left(-\frac{\phi_{,\varsigma_x\varsigma_x}(x)}{(\phi(x))^4} + \frac{4(\phi_{,\varsigma_x}(x))^4}{(\phi(x))^5} + \frac{2(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^4} \right) \nabla \rho \left(\frac{z}{\phi(x)} \right) \cdot z \\ &\quad \left. + \left(\frac{6(\phi_{,\varsigma_x}(x))^2}{(\phi(x))^4} - \frac{2\phi_{,\varsigma_x\varsigma_x}(x)}{(\phi(x))^3} \right) \rho \left(\frac{z}{\phi(x)} \right) \right| dz \\ &\leq c \int_{B_{\phi(x)}(0)} \frac{|u(x-z) - l_{\varsigma_x}(x-z)|}{(\phi(x))^4} dz (\|\nabla^2 \rho\|_\infty + \|\nabla \rho\|_\infty + \|\rho\|_\infty) \\ &\leq c \int_{B_{\phi(x)}(x)} |u(z) - l_{\varsigma_x}(z)| (\phi(x))^{-4} dz \\ (133) \quad &\leq c\epsilon^{-\frac{1}{2}}. \end{aligned} \quad (138)$$

Proof of Lemma completed. From Step 2, (126), for any $x \in N_\epsilon(\partial\Omega) \cap \Omega$ we have

$$\left| |\nabla\psi(x)|^2 - 1 \right|^2 \leq c\epsilon$$

so (112) follows. In the same way from Step 3, (113) follows.

Since for any $x \in \Omega \setminus N_\epsilon(\partial\Omega)$ we know $u(x) \geq \epsilon$ and so $\phi(x) = w(u(x)) = \epsilon$ and thus $\rho_{\phi(x)}(z) = \rho\left(\frac{z}{\epsilon}\right)\epsilon^{-1}$ and there for $\psi(x) = \int u(x-z)\rho_\epsilon(z)dz$. Thus (114) is established. $\square$

Lemma 9. *Let $\beta > 0$, suppose Ω is a convex set with $|\Omega \triangle B_1(0)| \leq \beta$. Let $u(z) = d(z, \partial\Omega)$. For any $x \in \Omega \setminus B_{\beta^{\frac{1}{8}}}(0)$ for which the approximate derivative ∇u exists*

$$\left| \nabla u(x) + \frac{x}{|x|} \right| \leq c\beta^{\frac{3}{16}}. \quad (139)$$

Proof. For any $x \in \Omega \setminus B_{\beta^{\frac{1}{8}}}(0)$ let $b_x \in \partial\Omega$ be such that $|b_x - x| = u(x)$. Recall $a_{\frac{x}{|x|}} = \partial\Omega \cap l_0^{\frac{x}{|x|}}$, we begin by showing

$$\left| b_x - \frac{x}{|x|} \right| \leq c\beta^{\frac{3}{16}}. \quad (140)$$

Using (110) from Lemma 6

$$|x - b_x| \leq \left| x - a_{\frac{x}{|x|}} \right| \stackrel{(110)}{\leq} 1 - |x| + c\sqrt{\beta}. \quad (141)$$

Hence

$$|x - b_x|^2 = |x|^2 - 2x \cdot b_x + |b_x|^2 \stackrel{(141)}{\leq} 1 - 2|x| + |x|^2 + c\sqrt{\beta}. \quad (142)$$

Therefor

$$\begin{aligned} -2x \cdot b_x &\stackrel{(142)}{\leq} 1 - 2|x| + c\sqrt{\beta} - |b_x|^2 \\ &\stackrel{(110)}{\leq} -2|x| + c\sqrt{\beta}. \end{aligned}$$

Thus $2|x| \leq 2x \cdot b_x + c\sqrt{\beta}$. Since $|x| > \beta^{\frac{1}{8}}$ we have

$$1 \leq \frac{x}{|x|} \cdot b_x + c\frac{\sqrt{\beta}}{|x|} \leq 1 + c\beta^{\frac{3}{8}}. \quad (143)$$

Hence

$$\left| b_x - \frac{x}{|x|} \right|^2 = |b_x|^2 + 1 - 2\frac{x}{|x|} \cdot b_x \stackrel{(143), (110)}{\leq} c\beta^{\frac{3}{8}}$$

which gives

$$\left| \frac{x}{|x|} - b_x \right| \leq c\beta^{\frac{3}{16}}. \quad (144)$$

Let $\theta_x = \frac{b_x}{|b_x|}$ so using Lemma 6 $\left| \eta_{b_x} + \frac{b_x}{|b_x|} \right| = |\eta_{a_{\theta_x}} + \theta_x| \stackrel{(111)}{\leq} c\beta^{\frac{1}{4}}$ and by (110) this easily implies

$$|\eta_{b_x} + b_x| \leq c\beta^{\frac{1}{4}}. \quad (145)$$

Now since $\nabla u(x) = \frac{x - b_x}{|x - b_x|} = \eta_{b_x}$ and so

$$\left| \nabla u(x) + \frac{x}{|x|} \right| \leq |\eta_{b_x} + b_x| + \left| \frac{x}{|x|} - b_x \right| \stackrel{(144), (145)}{\leq} c\beta^{\frac{3}{16}}$$

thus we have established (139). $\square$

Lemma 10. *Let Ω be a convex set and $|\Omega \triangle B_1| \leq \beta$. Define $u(x) = d(x, \partial\Omega)$, note that since u is convex ∇u is BV. Let $V(\nabla u, \cdot)$ denotes the total total variation of the measure ∇u . Firstly we have*

$$V(\nabla u, \Omega \setminus B_{3\beta^{\frac{1}{8}}}(0)) \leq c. \quad (146)$$

Secondly for any $\varepsilon \in (0, \beta^{\frac{1}{2}}]$, for any $x \in \Omega \setminus (N_{2\varepsilon}(\partial\Omega) \cup B_{4\beta^{\frac{1}{8}}}(0))$ we have

$$V(\nabla u, B_\varepsilon(x)) \leq c\beta^{\frac{3}{16}}\varepsilon. \quad (147)$$

Proof. Let $\tau \in (0, \varepsilon)$ be some small number. For any $x \in \Omega \setminus (N_{4\tau}(\partial\Omega) \cup B_{\frac{3}{2}\beta^{\frac{1}{8}}}(0)) =: \Pi_\tau$.

Let $w_\tau(x) = u * \rho_\tau(x)$ and $v^\tau = \frac{\nabla_x w_\tau}{|\nabla_x w_\tau|}$.

Note from Lemma 9 for any $x \in \Pi_\tau$

$$\begin{aligned} \left| \nabla w_\tau(x) + \frac{x}{|x|} \right| &= \left| \int \left(\nabla u(x-z) + \frac{x}{|x|} \right) \rho_\tau(z) dz \right| \\ &\leq \int \left| \left(\nabla u(x-z) + \frac{x-z}{|x-z|} \right) \rho_\tau(z) \right| dz + \int \left| \frac{x-z}{|x-z|} - \frac{x}{|x|} \right| \rho_\tau(z) dz \\ &\stackrel{(139)}{\leq} c \sup_{z \in B_\tau(0)} \left| \frac{x-z}{|x-z|} - \frac{x}{|x|} \right| + c\beta^{\frac{3}{16}} \\ &\leq c\beta^{\frac{3}{16}}. \end{aligned} \quad (148)$$

From this its easy to conclude that

$$\|w_\tau - \text{dist}(\cdot, \partial B_1(0))\|_{L^\infty(\Pi_\tau)} \leq c\beta^{\frac{3}{16}}. \quad (149)$$

Step 1. We will show that for any $t \in (8\tau, 1 - 2\beta^{\frac{1}{8}})$

$$\int_{w_\tau^{-1}(t)} |v_{1,1}^\tau(z) + v_{2,2}^\tau(z)| dH^1 z \leq c \quad (150)$$

Proof of Step 1. We define the ‘angle’ function by

$$A(x) := \begin{cases} \cos^{-1}\left(\frac{x_1}{|x|}\right) & \text{for } x_2 \geq 0 \\ 2\pi - \cos^{-1}\left(\frac{x_1}{|x|}\right) & \text{for } x_2 < 0 \end{cases} \quad (151)$$

Note that A is smooth except at the half line $\{(x_1, x_2) : x_2 = 0, x_1 > 1\}$. For $x \in \Pi_\tau$ we have $|v^\tau(x)|^2 = 1$, so

$$\partial_1(|v^\tau(x)|^2) = v_1^\tau(x)v_{1,1}^\tau(x) + v_2^\tau(x)v_{2,1}^\tau(x) = 0 \quad (152)$$

and

$$\partial_2(|v^\tau(x)|^2) = v_1^\tau(x)v_{1,2}^\tau(x) + v_2^\tau(x)v_{2,2}^\tau(x) = 0. \quad (153)$$

Since u is the 1-Lipschitz,

$$\|w_\tau - u\|_{L^\infty(\Pi_\tau)} \leq 2\tau, \quad (154)$$

and so from this and (149) we have that for any $t \in (8\tau, 1 - 2\beta^{\frac{1}{8}})$, $w_\tau^{-1}(t) \subset \Pi_\tau$ and hence v^τ is well defined along this level set. We also know that for any $x \in w_\tau^{-1}(t)$ the tangent to curve $w_\tau^{-1}(t)$ is given by $\begin{pmatrix} -v_2^\tau(x) \\ v_1^\tau(x) \end{pmatrix}$.

Now there exists a point $x_0 \in w_\tau^{-1}(t)$ such that $A\left(\frac{-v_2^\tau(x_0)}{v_1^\tau(x_0)}\right) = 0$. Let $\Phi^t : [0, H^1(w_\tau^{-1}(t))) \rightarrow w_\tau^{-1}(t)$ denote that parameterisation of Γ_t by arc-length with $\Phi^t(0) = x_0$, so $\dot{\Phi}^t(s) = \begin{pmatrix} -v_2^\tau(\Phi^t(s)) \\ v_1^\tau(\Phi^t(s)) \end{pmatrix}$.

Define $\Theta_t : [0, H^1(w_\tau^{-1}(t))] \rightarrow \mathbb{R}$ by $\Theta_t(s) = A(\dot{\Phi}^t(s))$. Now pick $s \in (0, H^1(w_\tau^{-1}(t)))$, suppose $v_1^\tau(\Phi^t(s)) > 0$, then

$$\begin{aligned}
\dot{\Theta}_t(s) &= \cos^{-1}(-v_2^\tau(\Phi^t(s))) \frac{\partial}{\partial t}(-v_2^\tau(\Phi^t(s))) \\
&= \cos^{-1}(-v_2^\tau(\Phi^t(s))) \left(-v_{2,1}^\tau(\Phi^t(s)) \dot{\Phi}_1^t(t) - v_{2,2}^\tau(\Phi^t(s)) \dot{\Phi}_2^t(t) \right) \\
&= \cos^{-1}(-v_2^\tau(\Phi^t(s))) (v_{2,1}^\tau(\Phi^t(s)) v_2^\tau(\Phi^t(s)) - v_{2,2}^\tau(\Phi^t(s)) v_1^\tau(\Phi^t(s))) \\
&\stackrel{(152)}{=} \cos^{-1}(-v_2^\tau(\Phi^t(s))) (-v_{1,1}^\tau(\Phi^t(s)) v_1^\tau(\Phi^t(s)) - v_{2,2}^\tau(\Phi^t(s)) v_1^\tau(\Phi^t(s))) \\
&= -\cos^{-1}(-v_2^\tau(\Phi^t(s))) v_1^\tau(\Phi^t(s)) (v_{1,1}^\tau(\Phi^t(s)) + v_{2,2}^\tau(\Phi^t(s))). \tag{155}
\end{aligned}$$

Now for any $w \in (-1, 1)$, $\cos^{-1}(w) = -(\sin(\cos^{-1}(w)))^{-1}$ so

$$\dot{\Theta}_t(t) = \frac{v_1^\tau(\Phi^t(s))}{\sin(\cos^{-1}(-v_2^\tau(\Phi^t(s))))} (v_{1,1}^\tau(\Phi^t(s)) + v_{2,2}^\tau(\Phi^t(s))). \tag{156}$$

Recall $\left| \frac{-v_2^\tau(\Phi^t(s))}{v_1^\tau(\Phi^t(s))} \right| = 1$ and we supposed $v_1^\tau(\Phi^t(s)) \geq 0$, so

$$\begin{aligned}
v_1^\tau(\Phi^t(s)) &= \sqrt{1 - (v_2^\tau(\Phi^t(s)))^2} \\
&= \sqrt{1 - (\cos(\cos^{-1}(-v_2^\tau(\Phi^t(s))))^2} \\
&= \sin(\cos^{-1}(-v_2^\tau(\Phi^t(s)))). \tag{157}
\end{aligned}$$

Thus from (156)

$$\dot{\Theta}_t(s) = (v_{1,1}^\tau(\Phi^t(s)) + v_{2,2}^\tau(\Phi^t(s))) \text{ for any } s \in (0, H^1(w_\tau^{-1}(t))) \text{ with } v_1^\tau(\Phi^t(s)) > 0. \tag{158}$$

Suppose we have $s \in (0, H^1(w_\tau^{-1}(t)))$ with $v_1^\tau(\Phi^t(s)) < 0$, then in the same way as (157) we have

$$v_1^\tau(\Phi^t(s)) = -\sqrt{1 - (\cos(\cos^{-1}(-v_2^\tau(\Phi^t(s))))^2} = -\sin(\cos^{-1}(-v_2^\tau(\Phi^t(s)))). \tag{159}$$

And since $v_1^\tau(\Phi^t(s)) < 0$, by definition of A (see (151)) arguing as in (156) we have

$$\begin{aligned}
\dot{\Theta}_t(s) &= \frac{-v_1^\tau(\Phi^t(s))}{\sin(\cos^{-1}(-v_2^\tau(\Phi^t(s))))} (v_{1,1}^\tau(\Phi^t(s)) + v_{2,2}^\tau(\Phi^t(s))) \\
&\stackrel{(159)}{=} v_{1,1}^\tau(\Phi^t(s)) + v_{2,2}^\tau(\Phi^t(s)) \text{ for } s \in (0, H^1(w_\tau^{-1}(t))) \text{ with } v_1^\tau(\Phi^t(s)) < 0.
\end{aligned}$$

Without loss of generality we can assume $|\{s \in [0, H^1(w_\tau^{-1}(t))] : v_1^\tau(\Phi^t(s)) = 0\}| = 0$. Thus by continuity of $\dot{\Theta}_t(\cdot)$, $v_{1,1}^\tau(\Phi^\tau(\cdot))$ and $v_{2,2}^\tau(\Phi^\tau(\cdot))$ we have

$$\dot{\Theta}_t(s) = v_{1,1}^\tau(\Phi^t(s)) + v_{2,2}^\tau(\Phi^t(s)) \text{ for } s \in [0, H^1(w_\tau^{-1}(t))). \tag{160}$$

Now since u is concave, w_τ is concave and so the set $w_\tau^{-1}([t, \infty))$ is a convex set, hence

$$v_{1,1}^\tau(\Phi^t(s)) + v_{2,2}^\tau(\Phi^t(s)) = \dot{\Theta}_t(s) \geq 0 \text{ for any } s \in [0, H^1(w_\tau^{-1}(t))). \tag{161}$$

Hence

$$\int_{w_\tau^{-1}(t)} |v_{1,1}^\tau(z) + v_{2,2}^\tau(z)| dH^1 z = \int_0^{H^1(w_\tau^{-1}(t))} \dot{\Theta}_t(s) ds \leq c$$

Step 2. Let $x \in \Pi_\tau$ and define

$$t_1 = \inf \{s \in \mathbb{R} : w_\tau^{-1}(s) \cap B_\varepsilon(x) \neq \emptyset\} \text{ and } t_2 = \sup \{s \in \mathbb{R} : w_\tau^{-1}(s) \cap B_\varepsilon(x) \neq \emptyset\}. \tag{162}$$

For any $t \in (t_1, t_2)$

$$\sup \{\Theta_t(s_1) - \Theta_t(s_2) : \Phi^t(s_1), \Phi^t(s_2) \in B_\varepsilon(x)\} \leq c\beta^{\frac{3}{10}}. \tag{163}$$

Proof of Step 2. Let $s_1, s_2 \in [0, H^1(w_\tau^{-1}(t))]$ such that $\Phi^t(s_1), \Phi^t(s_2) \in B_\varepsilon(x)$, since Φ^t is parameterization of Γ_t by arclength $\dot{\Phi}^t(s)$ is the unit tangent to $w_\tau^{-1}(t)$ at $\Phi^t(s)$. Thus

$$R\left(\frac{\nabla w_\tau(\Phi^t(s_i))}{|\nabla w_\tau(\Phi^t(s_i))|}\right) = \dot{\Phi}^t(s_i) \text{ for } i = 1, 2.$$

However by Lemma 9 (recalling the fact that $|\Phi^t(s_1)| > \frac{3\beta^{\frac{1}{8}}}{2}$ and $|\Phi^t(s_2)| > \frac{3\beta^{\frac{1}{8}}}{2}$ in order to apply the lemma)

$$\begin{aligned} |\nabla w_\tau(\Phi^t(s_1)) - \nabla w_\tau(\Phi^t(s_2))| &= \left| \int (\nabla u(\Phi^t(s_1) - z) - \nabla u(\Phi^t(s_2) - z)) \rho_\tau(z) dz \right| \\ &\stackrel{(139)}{\leq} c \int_{B_\tau(0)} \left| \frac{\Phi^t(s_1) - z}{|\Phi^t(s_1) - z|} - \frac{\Phi^t(s_2) - z}{|\Phi^t(s_2) - z|} \right| \rho_\tau(z) dz + c\beta^{\frac{3}{16}}. \end{aligned} \quad (164)$$

Note $z \in B_\tau(0) \subset B_{\beta^{\frac{1}{2}}}(0)$ so as $x \notin B_{2\beta^{\frac{1}{8}}}(0)$ we have $|\Phi^t(s_1) - z| \geq |\Phi^t(s_1)| - |z| \geq \beta^{\frac{1}{8}}$. Recall the elementary inequality

$$\left| \frac{z}{|z|} - \frac{y}{|y|} \right| \leq 2|z - y| \text{ for any } z, y \text{ with } |z| \geq 1, |y| \geq 1. \quad (165)$$

So in particular we have

$$\left\| \frac{\Phi^t(s_1) - z}{|\Phi^t(s_1) - z|} - \frac{\Phi^t(s_2) - z}{|\Phi^t(s_2) - z|} \right\| \leq \frac{2}{\beta^{\frac{1}{8}}} |\Phi^t(s_1) - \Phi^t(s_2)| \leq 2\beta^{\frac{3}{8}}. \quad (166)$$

So with (164) this gives

$$|\nabla w_\tau(\Phi^t(s_1)) - \nabla w_\tau(\Phi^t(s_2))| \leq c\beta^{\frac{3}{16}}. \quad (167)$$

Recall as a consequence of (148) we know

$$||\nabla w_\tau(x)| - 1| \leq c\beta^{\frac{3}{16}} \text{ for any } x \in \Pi_\tau \quad (168)$$

so

$$\begin{aligned} |\dot{\Phi}(s_1) - \dot{\Phi}(s_2)| &\stackrel{(167)}{\leq} \left| R\left(\frac{\nabla w_\tau(\Phi^t(s_1))}{|\nabla w_\tau(\Phi^t(s_1))|}\right) - R(\nabla w_\tau(\Phi^t(s_1))) \right| \\ &\quad + \left| R\left(\frac{\nabla w_\tau(\Phi^t(s_2))}{|\nabla w_\tau(\Phi^t(s_2))|}\right) - R(\nabla w_\tau(\Phi^t(s_2))) \right| + c\beta^{\frac{3}{16}} \\ &\stackrel{(168)}{\leq} c\beta^{\frac{3}{16}}. \end{aligned} \quad (169)$$

Thus

$$|\Theta_t(s_1) - \Theta_t(s_2)| = \left| A(\dot{\Phi}^t(s_1)) - A(\dot{\Phi}^t(s_2)) \right| \stackrel{(169)}{\leq} c\beta^{\frac{3}{16}} \quad (170)$$

and so (163) is established.

Proof of Lemma completed. For any $t \in (t_1, t_2)$. Let $s_1^t = \inf \{s : \Phi^t(s) \in B_\varepsilon(x)\}$, $s_2^t = \sup \{s : \Phi^t(s) \in B_\varepsilon(x)\}$. So $[s_1^t, s_2^t] = \{s : \Phi^t(s) \in B_\varepsilon(x)\}$. Now

$$\begin{aligned} \int_{[s_1^t, s_2^t]} |v_{1,1}^\tau(\Phi^t(s)) + v_{1,1}^\tau(\dot{\Phi}^t(s))| ds &\stackrel{(161)}{=} \int_{[s_1^t, s_2^t]} \dot{\Theta}_t(s) ds \\ &\stackrel{(170)}{\leq} c\beta^{\frac{3}{16}}. \end{aligned} \quad (171)$$

Thus

$$\begin{aligned}
\int_{B_\varepsilon(x)} |v_{1,1}^\tau(z) + v_{1,1}^\tau(z)| |\nabla w_\tau(z)| dz &= \int_{t_1}^{t_2} \int_{w_\tau^{-1}(t)} |v_{1,1}^\tau(z) + v_{1,1}^\tau(z)| dH^1 z dt \\
&= \int_{t_1}^{t_2} \int_{[s_1^t, s_2^t]} |v_{1,1}^\tau(\Phi^t(s)) + v_{1,1}^\tau(\Phi^t(s))| ds dt \\
&\stackrel{(171)}{\leq} c |t_1 - t_2| \beta^{\frac{3}{16}}.
\end{aligned}$$

By using (148) and recalling the definition (162) of Step 2 we must have $|t_1 - t_2| \leq c\varepsilon$. Also from (148) we know $|\nabla w_\tau(z)| \geq 1 - c\beta^{\frac{3}{16}}$ for all $z \in B_\varepsilon(x)$, so putting these things together we have

$$\int_{B_\varepsilon(x)} |v_{1,1}^\tau(z) + v_{1,1}^\tau(z)| dz \leq c\varepsilon \beta^{\frac{3}{16}}. \quad (172)$$

Since $v_{2,1}^\tau = v_{1,2}^\tau$ we have

$$\begin{pmatrix} v_{1,1}^\tau & v_{2,1}^\tau \\ v_{2,1}^\tau & v_{2,2}^\tau \end{pmatrix} \begin{pmatrix} v_1^\tau \\ v_2^\tau \end{pmatrix} \stackrel{(152),(153)}{=} \begin{pmatrix} 0 \\ 0 \end{pmatrix} \text{ and } \begin{pmatrix} v_{1,1}^\tau & v_{2,1}^\tau \\ v_{2,1}^\tau & v_{2,2}^\tau \end{pmatrix} \begin{pmatrix} -v_2^\tau \\ v_1^\tau \end{pmatrix} \stackrel{(152),(153)}{=} (v_{1,1}^\tau + v_{2,2}^\tau) \begin{pmatrix} -v_2^\tau \\ v_1^\tau \end{pmatrix}.$$

Letting $\|\cdot\|$ denote the operator norm of a matrix, since $\begin{pmatrix} v_1^\tau & -v_2^\tau \\ v_2^\tau & v_1^\tau \end{pmatrix} \in O(2)$ thus

$$\begin{aligned}
\left\| \begin{pmatrix} v_{1,1}^\tau & v_{2,1}^\tau \\ v_{2,1}^\tau & v_{2,2}^\tau \end{pmatrix} \right\| &= \left\| \begin{pmatrix} v_{1,1}^\tau & v_{2,1}^\tau \\ v_{2,1}^\tau & v_{2,2}^\tau \end{pmatrix} \begin{pmatrix} v_1^\tau & -v_2^\tau \\ v_2^\tau & v_1^\tau \end{pmatrix} \right\| \\
&= \left\| \begin{pmatrix} 0 & -(v_{1,1}^\tau + v_{2,2}^\tau)v_2^\tau \\ 0 & (v_{1,1}^\tau + v_{2,2}^\tau)v_1^\tau \end{pmatrix} \right\| \\
&\leq c |v_{1,1}^\tau + v_{2,2}^\tau|
\end{aligned}$$

As the operator norm and Euclidean norm are equivalent $|Dv^\tau| \leq c |v_{1,1}^\tau + v_{2,2}^\tau|$. Hence from (172)

$$\int_{B_\varepsilon(x)} |Dv^\tau| \leq c\beta^{\frac{3}{16}}\varepsilon. \quad (173)$$

Additionally for any $t \in (8\tau, 1 - 2\beta^{\frac{1}{8}})$, we have $\int_{w_\tau^{-1}(t)} |Dv^\tau| dH^1 z \stackrel{(150)}{\leq} c$. Note that by (149) and (154) we have $\Pi_{16\tau} \setminus B_{\frac{1}{3\beta^{\frac{1}{8}}}}(0) \subset w_\tau^{-1} \left([8\tau, 1 - 2\beta^{\frac{1}{8}}] \right)$ by using the Co-area formula

$$\int_{\Pi_{16\tau} \setminus B_{\frac{1}{3\beta^{\frac{1}{8}}}}(0)} |Dv^\tau| |\nabla w^\tau| \leq \int_{8\tau}^{1-2\beta^{\frac{1}{8}}} \int_{w_\tau^{-1}(s)} |Dv^\tau| dH^1 z ds \leq c.$$

Thus using (148)

$$\int_{\Pi_{8\tau} \setminus B_{\frac{1}{3\beta^{\frac{1}{8}}}}(0)} |Dv^\tau| \leq c \quad (174)$$

Let $\tau_n = 2^{-n}$. So given $x \in \Omega \setminus \left(N_{2\varepsilon}(\partial\Omega) \cup B_{\frac{1}{4\beta^{\frac{1}{8}}}}(0) \right)$ by definition of w_{τ_n} we know $\nabla w_{\tau_n} \xrightarrow{L^1} \nabla u$ and so

$$\begin{aligned}
\int_{B_\varepsilon(x)} |1 - |\nabla w_{\tau_n}|| &= \int_{B_\varepsilon(x)} ||\nabla u| - |\nabla w_{\tau_n}|| \\
&\leq \int_{B_\varepsilon(x)} |\nabla u - \nabla w_{\tau_n}| \rightarrow 0 \text{ as } n \rightarrow \infty.
\end{aligned}$$

So we have

$$\begin{aligned} \|v^{\tau_n} - \nabla u\|_{L^1(B_\varepsilon(x))} &\leq \| |\nabla w^{\tau_n}|^{-1} \nabla w_{\tau_n} - \nabla w_{\tau_n} \|_{L^1(B_\varepsilon(x))} + \|\nabla w_{\tau_n} - \nabla u\|_{L^1(B_\varepsilon(x))} \\ &\rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned} \quad (175)$$

Also as by (173) the sequence v^{τ_n} is bounded in BV and by (175) is converging in L^1 to u , by Proposition 3.13 [Am-Fu-Pa 00] we have that

$$v^{\tau_n} \xrightarrow{BV(B_\varepsilon(x))} \nabla u.$$

Hence by Remark 3.5 [Am-Fu-Pa 00] total variation is lower semicontinuous with respect to weak convergence we have $V(\nabla u, B_\varepsilon(x)) \leq c\beta^{\frac{3}{16}}\varepsilon$. By arguing in exactly the same way we see (174) implies $V(\Omega \setminus B_{\frac{1}{3\beta^{\frac{1}{8}}}}(0)) \leq c$. $\square$

Lemma 11. *Let Ω be a convex domain and $|\Omega \triangle B_1(0)| \leq \beta$. Let $u(x) = d(x, \partial\Omega)$ and for $\varepsilon \in (0, \frac{\beta^{\frac{1}{2}}}{4}]$ define $u_\varepsilon := u * \rho_\varepsilon$. For any a for which $a \in \Omega \setminus (N_{8\varepsilon}(\partial\Omega) \cup B_{\frac{1}{4\beta^{\frac{1}{8}}}}(0))$ we have*

$$|\nabla u_\varepsilon(x) - 1| \leq c\varepsilon^{-1}V(\nabla u, B_{4\varepsilon}(a)) \text{ for any } x \in B_{2\varepsilon}(a). \quad (176)$$

Proof. Firstly recall that since u is concave and hence ∇u is BV, so by Poincaré's inequality

$$\left(\int_{B_{4\varepsilon}(a)} |\nabla u - w|^2 \right)^{\frac{1}{2}} \leq cV(\nabla u, B_{4\varepsilon}(a)), \quad (177)$$

we have $w = \fint_{B_{4\varepsilon}(a)} \nabla u$. Using Holder's inequality we get another form of Poincaré's inequality

$$\int_{B_{4\varepsilon}(a)} |\nabla u - w| \leq c\|\nabla u - w\|_{L^2(B_{4\varepsilon}(a))}\varepsilon \leq c\varepsilon V(\nabla u, B_{4\varepsilon}(a)). \quad (178)$$

Now

$$\begin{aligned} \pi 16\varepsilon^2 |1 - |w|| &= \int_{B_{4\varepsilon}(a)} |1 - |w|| \\ &= \int_{B_{4\varepsilon}(a)} ||\nabla u| - |w|| \\ &\stackrel{(178)}{\leq} c\varepsilon V(\nabla u, B_{4\varepsilon}(a)). \end{aligned}$$

Thus $|1 - |w|| \leq c \frac{V(\nabla u, B_{4\varepsilon}(a))}{\varepsilon}$ and so there must exists $v \in S^1$ such that $|v - w| \leq |1 - |w||$ hence putting this together with (178) we have

$$\fint_{B_{4\varepsilon}(a)} |\nabla u - v| \leq c \frac{V(\nabla u, B_{4\varepsilon}(a))}{\varepsilon}. \quad (179)$$

Hence for any $w \in B_{2\varepsilon}(a)$, using Lemma 10 for the last inequality

$$\begin{aligned} |\nabla u_\varepsilon(w) - v| &= \left| \int (\nabla u(z) - v) \rho_\varepsilon(w - z) dz \right| \\ &\leq c\varepsilon^{-2} \left| \int (\nabla u(z) - v) \rho(\varepsilon^{-1}(z - w)) dz \right| \\ &\leq c\varepsilon^{-2} \int_{B_{2\varepsilon}(w)} |\nabla u(z) - v| dz \\ &\stackrel{(179)}{\leq} c \frac{V(\nabla u, B_{4\varepsilon}(a))}{\varepsilon}. \end{aligned} \quad (180)$$

This completes the proof of Lemma 11. $\square$

Lemma 12. *Let Ω be a convex domain and $|\Omega \triangle B_1(0)| \leq \beta$. Let $u(x) = d(x, \partial\Omega)$ and define $u_\varepsilon := u * \rho_\varepsilon$. Define $\Lambda := \Omega \setminus \left(N_{8\varepsilon}(\partial\Omega) \cup B_{\frac{1}{4\beta^{\frac{1}{8}}}}(0) \right)$, we will show that for any $\varepsilon \in (0, \frac{\beta^{\frac{1}{4}}}{4}]$*

$$\int_{\Lambda} \varepsilon^{-1} \left| 1 - |\nabla u_\varepsilon|^2 \right|^2 + \varepsilon |\nabla^2 u_\varepsilon|^2 \leq c\beta^{\frac{3}{16}}. \quad (181)$$

Proof of Lemma. By the 5r Covering Theorem ([Ma 95], Theorem 2.1) then we can find a finite collection of balls $J := \left\{ B_{\frac{2\varepsilon}{5}}(x_i) : i = 1, 2, \dots, m \right\}$ that are piecewise disjoint and $\Lambda \subset \bigcup_{i=1}^m B_{2\varepsilon}(x_i)$.

Note that for any $i = 1, 2, \dots, n$ since they are pairwise disjoint there are at most C_1 balls from J inside $B_{5\varepsilon}(x_i)$. Thus $\| \sum_{i=1}^m \mathbb{1}_{B_{5\varepsilon}(x_i)} \|_{L^\infty(\Omega)} \leq C_1$ and this obviously implies $\| \sum_{i=1}^m \mathbb{1}_{B_{2\varepsilon}(x_i)} \|_{L^\infty(\Omega)} \leq C_1$.

For $x, y \in \mathbb{R}^2$ let $x \otimes y := \begin{pmatrix} x_1 y_1 & x_1 y_2 \\ x_2 y_1 & x_2 y_2 \end{pmatrix}$. Now given $a \in \Lambda$ if $x \in B_{2\varepsilon}(a)$, let $w = f_{B_\varepsilon(x)} \nabla u$

$$\begin{aligned} |\nabla^2 u_\varepsilon(x)| &= \left| \int \nabla u(z) \otimes \nabla \rho_\varepsilon(x-z) dz \right| \\ &\leq \left| \int (\nabla u(z) - w) \otimes \nabla \rho \left(\frac{x-z}{\varepsilon} \right) \varepsilon^{-3} dz \right| \\ &\leq c\varepsilon^{-3} \left| \int_{B_\varepsilon(x)} (\nabla u - w) dz \right| \\ &\stackrel{(178)}{\leq} c\varepsilon^{-2} V(\nabla u, B_{4\varepsilon}(a)). \end{aligned} \quad (182)$$

So

$$\begin{aligned} \int_{\Lambda} |\nabla^2 u_\varepsilon|^2 dx &\leq \sum_{i=1}^m c \int_{B_{2\varepsilon}(x_i)} |\nabla^2 u_\varepsilon|^2 dx \\ &\leq c \sum_{i=1}^m \varepsilon^2 \|\nabla^2 u_\varepsilon\|_{L^\infty(B_{2\varepsilon}(x_i))}^2 \\ &\stackrel{(182)}{\leq} c\varepsilon^2 \left(\sum_{i=1}^m \varepsilon^{-4} (V(\nabla u, B_{4\varepsilon}(x_i)))^2 \right) \\ &\stackrel{(147)}{\leq} c\beta^{\frac{3}{16}} \varepsilon^{-1} \left(\sum_{i=1}^m V(\nabla u, B_{4\varepsilon}(x_i)) \right) \\ &\leq c\beta^{\frac{3}{16}} \varepsilon^{-1} V(\nabla u, \Lambda) \\ &\stackrel{(146)}{\leq} c\varepsilon^{-1} \beta^{\frac{3}{16}}. \end{aligned} \quad (183)$$

Now

$$\begin{aligned}
\int_{\Lambda} \left| 1 - |\nabla u_{\varepsilon}|^2 \right|^2 dz &\leq c \sum_{i=1}^m \int_{B_{2\varepsilon}(x_i)} |1 - |\nabla u_{\varepsilon}||^2 dz \\
&\stackrel{(147),(176)}{\leq} \sum_{i=1}^m c \varepsilon^2 \beta^{\frac{3}{16}} \| |1 - |\nabla u_{\varepsilon}|| \|_{L^{\infty}(B_{2\varepsilon}(x_i))} \\
&\stackrel{(176)}{\leq} \sum_{i=1}^m c \varepsilon \beta^{\frac{3}{16}} V(\nabla u, B_{4\varepsilon}(x_i)) \\
&\leq c \varepsilon \beta^{\frac{3}{16}} V(\nabla u, \Omega \setminus B_{2\beta^{\frac{1}{8}}}(0)) \\
&\stackrel{(146)}{\leq} c \beta^{\frac{3}{16}} \varepsilon.
\end{aligned} \tag{184}$$

Putting (184) together with (183) establishes (181). $\square$

Lemma 13. *Let $\eta(x) = |x|$, $\varepsilon > 0$ and define $\eta_{\varepsilon}(x) := \int \eta(z) \rho_{\varepsilon}(x - z) dz$. So*

$$\int_{B_1(0) \setminus B_{2\varepsilon}(0)} \left| 1 - |\nabla \eta_{\varepsilon}|^2 \right|^2 dz \leq c \log(\varepsilon^{-1}) \varepsilon^2 \tag{185}$$

and

$$\int_{B_1(0) \setminus B_{2\varepsilon}(0)} |\nabla^2 \eta_{\varepsilon}|^2 dz \leq c \log(\varepsilon^{-1}). \tag{186}$$

Proof of Lemma. Note for $x \notin B_{2\varepsilon}(0)$, $z \in B_{\varepsilon}(x)$

$$\begin{aligned}
\left| \frac{z}{|z|} - \frac{x}{|x|} \right| &\leq \left| \frac{z|x| - x|z|}{|z||x|} \right| \\
&\leq \left| \frac{z|x| - x|x|}{|z||x|} \right| + \left| \frac{x|x| - x|z|}{|z||x|} \right| \\
&\leq \frac{c\varepsilon}{|x| - \varepsilon}.
\end{aligned} \tag{187}$$

So for $x \notin B_{2\varepsilon}(0)$

$$\begin{aligned}
\left| \nabla \eta_{\varepsilon}(x) - \frac{x}{|x|} \right| &= \left| \int \rho_{\varepsilon}(x - z) \left(\frac{x}{|x|} - \frac{z}{|z|} \right) dz \right| \\
&\stackrel{(187)}{\leq} \frac{c\varepsilon}{|x| - \varepsilon}.
\end{aligned} \tag{188}$$

Since $\int \frac{x}{|x|} \otimes \nabla \rho_{\varepsilon}(x - z) dz = 0$, for any $x \notin B_{2\varepsilon}(0)$

$$\begin{aligned}
|\nabla^2 \eta_{\varepsilon}(x)| &= \left| \int \nabla \eta_{\varepsilon}(z) \otimes \nabla \rho_{\varepsilon}(x - z) dz \right| \\
&= \left| \int \left(\nabla \eta_{\varepsilon}(z) - \frac{z}{|z|} \right) \otimes \nabla \rho_{\varepsilon}(x - z) dz \right| + \left| \int \left(\frac{x}{|x|} - \frac{z}{|z|} \right) \otimes \nabla \rho_{\varepsilon}(x - z) dz \right| \\
&\stackrel{(187),(188)}{\leq} \frac{c\varepsilon}{|x| - \varepsilon} \left| \int \nabla \rho_{\varepsilon}(x - z) dz \right| \\
&\leq \frac{c}{|x| - \varepsilon}.
\end{aligned} \tag{189}$$

Hence

$$\begin{aligned}
\int_{B_1(0) \setminus B_{2\varepsilon}(0)} |\nabla^2 \eta_\varepsilon(x)|^2 dx &= c \int_{2\varepsilon}^1 \int_{\partial B_h(0)} \left(\frac{1}{|z| - \varepsilon} \right)^2 dH^1 z dr \\
&\leq c \int_{\varepsilon}^1 \frac{1}{r} dr \\
&\leq c \log(\varepsilon^{-1})
\end{aligned} \tag{190}$$

which establish (186).

Note $||\nabla \eta_\varepsilon(x)| - 1|^2 \leq \left| \nabla \eta_\varepsilon(x) - \frac{x}{|x|} \right|^2 \stackrel{(188)}{\leq} c \frac{\varepsilon^2}{(|x| - \varepsilon)^2}$ so arguing in the same way as in (190) we have (185). $\square$

Lemma 14. *Let Ω be a convex domain and $|\Omega \triangle B_1(0)| \leq \beta$.*

Let $u(x) = d(x, \partial\Omega)$ and $\eta(x) = 1 - 10\beta^{\frac{3}{32}} + |x|$. Define $\Gamma := \{x : u(x) = \eta(x)\}$, we will show Γ is the boundary of a convex set with $H^1(\Gamma) \leq c\beta^{\frac{3}{32}}$,

$$\Gamma \subset N_{c\beta^{\frac{3}{16}}}(\partial B_{5\beta^{\frac{3}{32}}}(0)) \tag{191}$$

and for any $\varepsilon \in (0, \beta^{\frac{3}{16}}]$

$$|N_{2\varepsilon}(\Gamma)| \leq c\varepsilon\beta^{\frac{3}{32}}. \tag{192}$$

Proof of Lemma.

Step 1. We will show $\Pi := \{x \in \Omega : \eta(x) \leq u(x)\}$ is convex.

Proof of Step 1. Take $a, b \in \Pi$ and pick $\lambda \in [0, 1]$. Since u is concave $u(\lambda a + (1 - \lambda)b) \geq \lambda u(a) + (1 - \lambda)u(b)$ and since η is convex $\eta(\lambda a + (1 - \lambda)b) \leq \lambda \eta(a) + (1 - \lambda)\eta(b)$. Hence as $a, b \in \Pi$, $u(\lambda a + (1 - \lambda)b) \geq \eta(\lambda a + (1 - \lambda)b)$. Thus $[a, b] \subset \Pi$ and thus the set Π is convex.

Step 2. We will establish (191).

Proof of Step 2. Let $x \in \Gamma$ and let $b_x \in \partial\Omega$ be such that $|x - b_x| = u(x)$. So

$$1 - 10\beta^{\frac{3}{32}} + |x| = |b_x - x|. \tag{193}$$

And thus $1 - 10\beta^{\frac{3}{32}} + |x| \geq |b_x| - |x|$, so using (110)

$$2|x| \geq |b_x| - 1 + 10\beta^{\frac{3}{32}} \geq 10\beta^{\frac{3}{32}} - c\sqrt{\beta}. \tag{194}$$

Also from (193) we have

$$|x| = |b_x - x| - (1 - 10\beta^{\frac{3}{32}}) \stackrel{(110)}{\leq} 10\beta^{\frac{3}{32}} + \sqrt{\beta}. \tag{195}$$

Now using Lemma 9, since $\nabla u(x) = \frac{x - b_x}{|x - b_x|}$ so

$$\begin{aligned}
\left| \frac{x}{|x|} - \frac{b_x}{|b_x|} \right| &\leq \left| \frac{b_x - x}{|b_x - x|} - \frac{b_x}{|b_x|} \right| + \left| \frac{x - b_x}{|x - b_x|} + \frac{x}{|x|} \right| \\
&\stackrel{(195), (139)}{\leq} c\beta^{\frac{3}{32}}
\end{aligned} \tag{196}$$

so

$$\left| 1 - \frac{b_x}{|b_x|} \cdot \frac{x}{|x|} \right| = 2^{-1} \left| \frac{b_x}{|b_x|} - \frac{x}{|x|} \right|^2 \leq c\beta^{\frac{3}{16}}. \tag{197}$$

Again by Lemma 9 we have

$$\begin{aligned}
\left| |b_x - x| + (x - b_x) \cdot \frac{x}{|x|} \right| &\leq |x - b_x| \left| \frac{x - b_x}{|x - b_x|} + \frac{x}{|x|} \right| \\
&\leq 2 \left| \nabla u(x) + \frac{x}{|x|} \right| \\
&\stackrel{(139)}{\leq} c\beta^{\frac{3}{16}}
\end{aligned} \tag{198}$$

and thus

$$\begin{aligned}
\left| 2x \cdot \frac{x}{|x|} - 10\beta^{\frac{3}{32}} \right| &\stackrel{(197)}{\leq} \left| -10\beta^{\frac{3}{32}} + 1 - \frac{b_x}{|b_x|} \cdot \frac{x}{|x|} + 2x \cdot \frac{x}{|x|} \right| + c\beta^{\frac{3}{16}} \\
&= \left| 1 - 10\beta^{\frac{3}{32}} + |x| - \left(\frac{b_x}{|b_x|} - x \right) \cdot \frac{x}{|x|} \right| + c\beta^{\frac{3}{16}} \\
&\stackrel{(193)}{=} \left| |b_x - x| - \left(\frac{b_x}{|b_x|} - x \right) \cdot \frac{x}{|x|} \right| + c\beta^{\frac{3}{16}} \\
&\leq \left| |b_x - x| + (x - b_x) \cdot \frac{x}{|x|} \right| + c\beta^{\frac{3}{16}} \\
&\stackrel{(198)}{\leq} c\beta^{\frac{3}{16}}
\end{aligned}$$

hence $\left| 2|x| - 10\beta^{\frac{3}{32}} \right| \leq c\beta^{\frac{3}{16}}$ for any $x \in \Gamma$, so (191) is established.

Since (191) implies the diameter of Π is bounded by $c\beta^{\frac{3}{32}}$ and since Π is a convex set it follows immediately that $H^1(\Gamma) \leq c\beta^{\frac{3}{32}}$.

Now the set Γ equipped with the Euclidean norm is a boundly compact metric space. So by applying the 5r Covering Theorem (Theorem 2.1 [Ma 95]) we can find a disjoint collection of balls $B_{2\varepsilon}(x_1), B_{2\varepsilon}(x_2), \dots, B_{2\varepsilon}(x_M)$ with $x_1, x_2, \dots, x_M \in \Gamma$ such that $\partial\Omega \subset \bigcup_{i=1}^M B_{10\varepsilon}(x_i)$. This implies $N_{2\varepsilon}(\partial\Omega) \subset \bigcup_{i=1}^M B_{20\varepsilon}(x_i)$. Since $H^1(\Gamma) \leq c\beta^{\frac{3}{32}}$ so $M \leq c\varepsilon^{-1}\beta^{\frac{3}{32}}$ and thus $|N_{2\varepsilon}(\Gamma)| \leq c\varepsilon\beta^{\frac{3}{32}}$ which establishes (192). $\square$

4.2. Proof of Proposition 1. Let $u(x) = d(x, \partial\Omega)$, let $w : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be the smooth monotonic function from the proof of Lemma 8, so w satisfies (116) and $\sup |\ddot{w}| \leq c\varepsilon^{-1}$ as in Lemma 8 for $x \in N_\varepsilon(\partial\Omega) \cap \Omega$ define $\phi(x) = w(u(x))$. Let

$$v(x) := \min \left\{ u(x), 1 - 10\beta^{\frac{3}{32}} + |x| \right\}. \tag{199}$$

and define

$$\xi(x) = \int v(x - z) \rho_{\phi(x)}(z) dz. \tag{200}$$

Let $\Pi = \left\{ x : u(x) > 1 - 10\beta^{\frac{3}{32}} + |x| \right\}$, and define $\Lambda_0 = \Omega \setminus (N_\varepsilon(\partial\Omega) \cup N_\varepsilon(\Pi))$, note that $\xi(x) = u_\varepsilon(x)$ for any $x \in \Lambda_0$.

Recall the function ψ defined in Lemma 8, note that $\xi(x) = \psi(x)$ for any $x \in N_\varepsilon(\partial\Omega) \cap \Omega$ thus

$$\int_{N_\varepsilon(\partial\Omega) \cap \Omega} \varepsilon^{-1} \left| 1 - |\nabla \xi|^2 \right|^2 + \varepsilon |\nabla^2 \xi|^2 \stackrel{(112), (113)}{\leq} c\varepsilon$$

Since $\psi = u_\varepsilon$ in Λ_0 , from (181) we have $\int_{\Lambda_0} \varepsilon^{-1} \left| 1 - |\nabla \xi|^2 \right|^2 + \varepsilon |\nabla^2 \xi|^2 \leq c\beta^{\frac{3}{16}}$ and so putting this two inequalities together we have

$$\int_{\Omega \setminus N_\varepsilon(\Pi)} \varepsilon^{-1} \left| 1 - |\nabla \xi|^2 \right|^2 + \varepsilon |\nabla^2 \xi|^2 \leq c\beta^{\frac{3}{16}} \tag{201}$$

Now as for any $x \in \Pi \setminus N_\epsilon(\partial\Pi)$, $w(x) = 1 - 10\beta^{\frac{3}{32}} + |x|$ and so $u_\epsilon(x) = \eta_\epsilon(x) + (1 - 10\beta^{\frac{3}{32}})$ where $\eta(x) = |x|$ and $\eta_\epsilon = \eta * \rho_\epsilon$. So $\nabla \xi(x) = \nabla \eta_\epsilon(x)$ and $\nabla^2 \xi(x) = \nabla^2 \eta_\epsilon(x)$ thus applying Lemma 13 we have

$$\int_{\Pi \setminus N_\epsilon(\partial\Pi)} \epsilon^{-1} \left| 1 - |\nabla \xi|^2 \right|^2 + \epsilon |\nabla^2 \xi|^2 \stackrel{(185), (186)}{\leq} c\epsilon \log(\epsilon^{-1}). \quad (202)$$

Since v is Lipschitz, so ξ is Lipschitz and so from (192) we have

$$\int_{N_\epsilon(\partial\Pi)} \epsilon^{-1} \left| 1 - |\nabla \xi|^2 \right|^2 \leq c\beta^{\frac{3}{32}}. \quad (203)$$

And note for any $x \in \Omega \setminus N_\epsilon(\partial\Omega)$

$$|\nabla^2 \xi(x)| = \epsilon^{-3} \left| \int \nabla v(z) \cdot \nabla \rho\left(\frac{x-z}{\epsilon}\right) dz \right| \leq c\epsilon^{-1}$$

so

$$\begin{aligned} \int_{N_\epsilon(\partial\Pi)} \epsilon |\nabla^2 \xi|^2 &\leq c\epsilon^{-1} |N_\epsilon(\partial\Pi)| \\ &\stackrel{(192)}{\leq} c\beta^{\frac{3}{32}}. \end{aligned} \quad (204)$$

Putting these inequalities together we have

$$\int_{N_\epsilon(\partial\Pi)} \epsilon^{-1} \left| 1 - |\nabla \xi|^2 \right|^2 + \epsilon |\nabla^2 \xi|^2 \leq c\beta^{\frac{3}{32}}. \quad (205)$$

Now inequalities (201), (202) and (205) give us that ξ satisfies (107). And since $\xi(x) = \psi(x)$ on $N_\epsilon(\partial\Omega) \cap \Omega$ from (115) satisfies $\nabla \xi(x) \cdot \eta_x = 1$ for any $x \in \partial\Omega$. This completes the proof of Proposition 1. $\square$

4.3. Proof Corollary 1. Let $\alpha = \inf_y |\Omega \triangle B_1(y)|$, without loss of generality assume $|\Omega \triangle B_1(0)| = \alpha$. Let $\beta = 4(\alpha + \epsilon)$, note that $\epsilon \in (0, \frac{\beta^{\frac{1}{2}}}{4})$ and $|\Omega \triangle B_1(0)| \leq \beta$.

Since $\xi \in \Lambda(\Omega)$ from (107) we have that $\inf_{u \in \Lambda(\Omega)} I_\epsilon(u) \leq c\beta^{\frac{3}{16}}$. Let $v \in \Lambda(\Omega)$ be the minimiser of I_ϵ and since v satisfies

$$\int_{\Omega} \left| 1 - |\nabla v|^2 \right| |\nabla^2 v| dz \leq \int_{\Omega} \epsilon^{-1} \left| 1 - |\nabla v|^2 \right|^2 + \epsilon |\nabla^2 v|^2 \leq c\beta^{\frac{3}{32}}$$

and as $\epsilon \in (0, \frac{\beta^{\frac{1}{2}}}{4}]$

$$\int_{\Omega} \left| 1 - |\nabla v|^2 \right|^2 dz \leq c\beta^{\frac{11}{16}}. \quad (206)$$

So defining $\tilde{\beta} = \beta^{\frac{3}{32}}$, we have that (6), (7) are satisfied and hence by Theorem 2

$$\int_{\Omega} \left| \nabla v(z) + \frac{z}{|z|} \right|^2 dz \leq c\tilde{\beta}^{\frac{1}{256}} = c\beta^{\frac{3}{32 \times 256}}.$$

Applying Lemma 9 we have

$$\int_{\Omega \setminus B_{\frac{1}{8}}(0)} |\nabla v - \nabla \zeta|^2 \leq c\beta^{\frac{3}{32 \times 256}}. \quad (207)$$

Now

$$\begin{aligned}
 \int_{B_{\beta^{\frac{1}{8}}}(0)} |\nabla v - \nabla \zeta|^2 dz &\leq \int_{B_{\beta^{\frac{1}{8}}}(0)} |\nabla v|^2 + 2|\nabla v| + 1 dz \\
 &\leq \int_{B_{\beta^{\frac{1}{8}}}(0)} \left(|1 - |\nabla v|^2| + c \right) dz \\
 &\stackrel{(206)}{\leq} c\beta^{\frac{1}{4}}
 \end{aligned}$$

together with (207) this gives $\|v - \zeta\|_{W^{2,2}(\Omega)} \leq c\beta^{\frac{3}{8192}} \leq c(\epsilon + \alpha)^{\frac{3}{8192}}$. $\square$

REFERENCES

- [Al-Ri-Se 02] F. Alouges; T. Riviere; S. Serfaty. Neel and cross-tie wall energies for planar micromagnetic configurations. A tribute to J. L. Lions. ESAIM Control Optim. Calc. Var. 8 (2002), 31–68
- [Am-De-Ma 99] L. Ambrosio; C. De Lellis; C. Mantegazza, Line energies for gradient vector fields in the plane. Calc. Var. Partial Differential Equations 9 (1999), no. 4, 327–255.
- [Am-Le-Ri 03] L. Ambrosio; M. Lecumberry; T. Riviere A viscosity property of minimizing micromagnetic configurations. Comm. Pure Appl. Math. 56 (2003), no. 6, 681–688
- [Am-Ki-Ri 02] L. Ambrosio; B. Kirchheim; M. Lecumberry; T. Riviere. On the rectifiability of defect measures arising in a micromagnetics model. Nonlinear problems in mathematical physics and related topics, II, 29–60, Int. Math. Ser. (N. Y.), 2, Kluwer/Plenum, New York, 2002.
- [Av-Gi 87] P. Aviles; Y. Giga, A mathematical problem related to the physical theory of liquid crystal configurations. Miniconference on geometry and partial differential equations. Proc. Centre Math. Anal. Austral. Nat. Univ., 12, Austral. Nat. Univ., Canberra, 1987.
- [Av-Gi 96] P. Aviles; Y. Giga, On lower semicontinuity of a defect energy obtained by a singular limit of the Ginzburg-Landau type energy for gradient fields. Proc. Roy. Soc. Edinburgh Sect. A 129 (1999), no. 1, 1–17.
- [Am-Fu-Pa 00] L. Ambrosio; N. Fusco; D. Pallara. Functions of bounded variation and free discontinuity problems. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000.
- [De-Ot 03] C. De Lellis, F. Otto, Structure of entropy solutions to the eikonal equation. J. Eur. Math. Soc. (JEMS) 5 (2003), no. 2, 107–145.
- [De-Ot-We 03] C. De Lellis, F. Otto, Felix; Westdickenberg, Michael Structure of entropy solutions for multi-dimensional scalar conservation laws. Arch. Ration. Mech. Anal. 170 (2003), no. 2, 137–184
- [De-Ko-Mu-Ot 00] A. DeSimone; S. Muller, R. Kohn, F. Otto. A compactness result in the gradient theory of phase transitions. Proc. Roy. Soc. Edinburgh Sect. A 131 (2001), no. 4, 833–844.
- [Di-Li-Me 91] R. DiPerna; P. Lions; Y. Meyer. L^p regularity of velocity averages. Ann. Inst. H. Poincaré Anal. Non Linéaire 8 (1991), no. 3-4, 271–287.
- [Ev-Ga 92] L.C. Evans. R.F. Gariepy. Measure theory and fine properties of functions. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1992.
- [Fed 69] H. Federer. Geometric measure theory. Die Grundlehren der mathematischen Wissenschaften, Band 153 Springer-Verlag New York Inc., New York 1969.
- [Fo-Ga 95] I. Fonseca, W. Gangbo. Degree theory in analysis and applications. Oxford Lecture Series in Mathematics and its Applications, 2. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1995.
- [Gi-Or 94] G. Gioia, M. Ortiz. The morphology and folding patterns of buckling-driven thin-film blisters. J. Mech. Phys. Solids 42 (1994), no. 3, 531–559.
- [Gi-Or 97] G. Gioia, M. Ortiz. Delamination of compressed thin films. Adv. Appl. Mech. 33 (1997) 119–192.
- [Ja-Pe 97] P. Jabin; B. Perthame. Compactness in Ginzburg-Landau energy by kinetic averaging. Comm. Pure Appl. Math. 54 (2001), no. 9, 1096–1109.
- [Ja-Ot-Pe 02] P. Jabin; F. Otto; B. Perthame. Line-energy Ginzburg-Landau models: zero-energy states. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) (2002), no. 1, 187–202.
- [Ji-Ko 00] W. Jin, R.V. Kohn, Singular perturbation and the energy of folds. J. Nonlinear Sci. 10 (2000), no. 3, 355–390.
- [Le-Ri 02] M. Lecumberry; T. Riviere. Regularity for micromagnetic configurations having zero jump energy. Calc. Var. Partial Differential Equations 15 (2002), no. 3, 389–402.
- [Lo pr] A. Lorent. A Poincaré type inequality for Aviles Giga energy and applications. Preprint.
- [Ma 95] P. Mattila. Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability. Cambridge Studies in Advanced Mathematics, 44. Cambridge University Press, Cambridge, 1995.

- [Mo-Mo 00] L. Modica, S. Mortola. Un esempio di Γ -convergenza. Boll. Un. Mat. Ital. B (5) 14 (1977), no. 1, 285–299.
- [Mu 78] F. Murat. Compacite par compensation: condition necessaire et suffisante de continuite faible sous une hypothese de rang constant. (French) Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 8 (1981), no. 1, 69–102.
- [Ri-Se 01] T. Riviere, S. Serfaty. Limiting domain wall energy for a problem related to micromagnetics. Comm. Pure Appl. Math. 54 (2001), no. 3, 294–338.
- [Ta 79] Tartar, L. Compensated compactness and applications to partial differential equations. Nonlinear analysis and mechanics: Heriot-Watt Symposium, Vol. IV, pp. 136–212, Res. Notes in Math., 39, Pitman, Boston, Mass.-London, 1979. 136–212.

MATHEMATICS DEPARTMENT, UNIVERSITY OF CINCINNATI, 2600 CLIFTON AVE., CINCINNATI, OHIO 45221
E-mail address: `lorentaw@uc.edu`